

MODEL THEORY AND THE TANNAKIAN FORMALISM

MOSHE KAMENSKY

ABSTRACT. We draw the connection between the model theoretic notions of internality and the binding group on one hand, and the Tannakian formalism on the other.

More precisely, we deduce the fundamental results of the Tannakian formalism by associating to a Tannakian category a first order theory, and applying the results on internality there. In the other direction, we formulate prove a general categorical statement that can be viewed as a “non-linear” version of the Tannakian formalism, and deduce the model theoretic result from it. For dessert, we formulate a version of the Tannakian formalism for differential linear groups, and show how the same techniques can be used to deduce the analogous results in that context.

INTRODUCTION

The aim of this paper is to exhibit the analogy and relationship between two seemingly unrelated theories. On the one hand, the Tannakian formalism, giving a duality theory between affine group schemes (or, more generally, gerbs) and a certain type of categories with additional structure, the Tannakian categories. On the other hand, a general notion of internality in model theory, valid for an arbitrary first order theory, that gives rise to a definable Galois group. The analogy is made precise by deriving (a weak version of) the fundamental theorem of the Tannakian duality (3.7) using the model theoretic internality.

The Tannakian formalism assigns to a group $\mathcal{G}$ over a field $\mathbb{k}$, its category of representations $\mathcal{R}ep_{\mathcal{G}}$. In the version we are mainly interested in, due to Saavedra (Saavedra-Rivano [28]), the group is an affine group scheme over a field. A similar approach works with groups in other categories, the first due to Krein and Tannaka, concerned with locally compact topological groups. Another example is provided in section 5. In the algebraic case, the category is the category of algebraic finite-dimensional representations. This is a $\mathbb{k}$ -linear category, but the category structure alone is not sufficient to recover the group. One therefore considers the additional structure given by the tensor product. The Tannakian formalism says that $\mathcal{G}$ can be recovered from this structure, together with the forgetful functor to the category of vector-spaces. The other half of the theory is a description of the tensor categories that arise as categories of representations: any tensor category satisfying suitable axioms is of the form $\mathcal{R}ep_{\mathcal{G}}$, provided it has a “fibre functor” into the category of vector spaces. Our main references for this subject are the first three sections of Deligne and Milne [5] and Deligne [4].

In model theory, internality was discovered by Zilber as a tool to study the structure of strongly minimal structure (Zil’ber [31]). Later, Poizat realised (in Poizat [27]) that this notion can be used to treat the Galois theory of differential equations. The definable Galois correspondence outlined in 2.5 has its origins there. Later,

the theory was generalised to larger classes of theories (Hrushovski [13], Hart and Shami [10], etc.), and applied in various contexts (e.g., Pillay [25] extended the differential Galois theory to arbitrary “ D -groups” definable in DCF).

In appendix B of Hrushovski [11], internality was reformulated in a way that holds in an arbitrary theory. One is interested in the group of automorphisms G of a definable set $\mathbf{Q}$ over another definable set $\mathbf{C}$. $\mathbf{Q}$ is internal to another set $\mathbf{C}$ if, after extending the base parameters, any element of $\mathbf{Q}$ is definable over the elements of $\mathbf{C}$. The idea is that under this condition, $\mathbf{Q}$ is close enough to $\mathbf{C}$ so that the G has a chance to be definable, but the requirement that a base extension is required prevents it from being trivial. The theorem is that indeed G is the group of points of a (pro-) definable group (see 2.5). This theory is reformulated again in Hrushovski [12], where the same construction is described as an abstract duality theory between definable groupoids in a theory $\mathcal{T}$, and certain expansions of it, called internal covers. It is this formulation that we use.

In section 4 we apply this theory by constructing, for a tensor category $\mathcal{C}$ an internal cover $T_{\mathcal{C}}$ of $ACF_{\mathbb{k}}$. Models of this theory correspond, roughly, with fibre functors on $\mathcal{C}$. The theory of internality provides a definable group in $ACF_{\mathbb{k}}$, and this is the group corresponding to $\mathcal{C}$. The other parts of the theory follow from the Galois theory, and from the abstract duality theory of Hrushovski [12]. The proof has the advantage that it is simple and more “geometric” than the algebraic one (though the result is slightly weaker). In section 5 we then define differential tensor categories, and explain how precisely the same approach gives an analogous theorem there. A similar result, using a somewhat different language was first obtain using algebraic methods in Ovchinnikov [23, 24]. It seems obvious that similar formalisms are possible in other contexts (e.g., difference fields, real closed fields).

The fundamental definitions and results required for the paper are explain in the first three sections. More precisely, the definitions and properties of tensor categories are given in sections 1 and 3. In section 2 we give some model theoretic background, including the statement of the internality theorem. In the first two sections we also explain how we can “go back” and deduce (again, a weak version of) the internality theorem from a categorical statement, which can be viewed as an abstract version of the Tannakian formalism. The categorical statement is in 1.13, and its application to internality starts with the translation between the languages in 2.9.

0.1. Questions. (Probably best read after the rest of the paper). Several questions (some vague) remain unanswered in the paper. The main ones, from my point of view, are the following.

0.1.1. As described in 4.12 and 4.13, the main results of Deligne [4] have very natural model theoretic translations. It seems reasonable to expect that there is a model theoretic proof, especially of the main theorem (which translates to having no new structure on $ACF_{\mathbb{k}}$), but I could not find it. The case when $\mathcal{C}$ is neutral appears to be easier from the algebraic point of view (and is proven already in Deligne and Milne [5]), but model theoretically I don’t know how to do even this case.

0.1.2. Conversely, the results and methods of Deligne and Milne [5] and Deligne [4] seem to suggest that one can do “model theory” inside a Tannakian category (and perhaps more generally in a tensor category). For example, as can be seen from 4.10, the statement of 3.7.2 can be viewed as saying that $\mathcal{C}$ has elimination of

quantifiers and elimination of imaginaries. I think it would be interesting to find out if this makes sense, and make it precise.

0.1.3. What is a general model theoretic machinery to prove results as in 4.7, but in full generality? (rather than just for fields). This can also be asked purely in terms of functors (though in this context there is probably no answer): Assume there is a functor $\mathcal{F}$ from $\mathbb{k}$ -algebras to sets (or groups), and a scheme (or group scheme) $\mathcal{G}$ with a map $\mathcal{G} \rightarrow \mathcal{F}$ that is a bijection on fields. What are conditions on $\mathcal{F}$ that allow us to deduce that this map is an isomorphism?

0.2. **Acknowledgements.** I would like to thank Ehud Hrushovski for suggesting this approach, and for help along the way. I would also like to thank Anand Pillay for his interest and useful remarks.

1. MONOIDAL CATEGORIES

We start with an overview of some of the notions related to monoidal categories. See Mac Lane [17, Ch. VII] or Deligne and Milne [5, sec. 1] for details (but note that we use the term “symmetric monoidal category” instead of “tensor category”; we use the latter for the abelian case).

1.1. **Monoidal categories.** A *symmetric monoidal category* is given by a tuple $(\mathcal{C}, \otimes, \phi, \psi)$, where:

- (1) $\mathcal{C}$ is a category
- (2) $\otimes$ is a functor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, $(\mathbf{X}, \mathbf{Y}) \mapsto \mathbf{X} \otimes \mathbf{Y}$ (in other words, the operation is functorial in each coordinate separately).
- (3) ϕ is a collection of functorial isomorphisms

$$\phi_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}} : (\mathbf{X} \otimes \mathbf{Y}) \otimes \mathbf{Z} \xrightarrow{\sim} \mathbf{X} \otimes (\mathbf{Y} \otimes \mathbf{Z})$$

one for each triple $\mathbf{X}, \mathbf{Y}, \mathbf{Z}$ of objects of $\mathcal{C}$. It is called the *associativity constraint*.

- (4) ψ is a collection of functorial isomorphisms $\psi_{\mathbf{X}, \mathbf{Y}} : \mathbf{X} \otimes \mathbf{Y} \xrightarrow{\sim} \mathbf{Y} \otimes \mathbf{X}$, called the *commutativity constraint*.

The commutativity constraint is required to satisfy $\psi_{\mathbf{X}, \mathbf{Y}} \circ \psi_{\mathbf{Y}, \mathbf{X}} = id_{\mathbf{X} \otimes \mathbf{Y}}$ for all objects $\mathbf{X}, \mathbf{Y}$. In addition, ϕ and ψ are required to satisfy certain “pentagon” and “hexagon” identities, which ensure that any two tensor expressions computed from the same set of objects are *canonically* isomorphic.

Finally, $\mathcal{C}$ is required have an *identity object*: this is an object $\mathbf{1}$, together with an isomorphism $u : \mathbf{1} \rightarrow \mathbf{1} \otimes \mathbf{1}$, such that $\mathbf{X} \mapsto \mathbf{1} \otimes \mathbf{X}$ is an equivalence of categories. It follows (Deligne and Milne [5, prop. 1.3]) that $(\mathbf{1}, u)$ is unique up to a unique isomorphism with the property that u can be uniquely extended to isomorphisms $l_{\mathbf{X}} : \mathbf{X} \rightarrow \mathbf{1} \otimes \mathbf{X}$, commuting with ϕ and ψ .

We denote by $\Gamma(\mathbf{X})$ (*global sections* of $\mathbf{X}$) the set $Hom(\mathbf{1}, \mathbf{X})$. Using the map u , we get a map $\Gamma(\mathbf{X}) \times \Gamma(\mathbf{Y}) \rightarrow \Gamma(\mathbf{X} \otimes \mathbf{Y})$, $(f, g) \mapsto (f \otimes g) \circ u$.

1.2. **Monoidal functors.** Let $\mathcal{C}$ and $\mathcal{D}$ be symmetric monoidal categories. A *monoidal functor* from $\mathcal{C}$ to $\mathcal{D}$ is a pair (F, χ) , where $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, and χ is a collection of isomorphisms $\chi_{\mathbf{X}, \mathbf{Y}} : F(\mathbf{X}) \otimes F(\mathbf{Y}) \xrightarrow{\sim} F(\mathbf{X} \otimes \mathbf{Y})$, compatible with the constraints, such that (F, χ) takes identity objects to identity objects.

If (F, χ) and (G, δ) are two tensor functors from $\mathcal{C}$ to $\mathcal{D}$, a map from (F, χ) to (G, δ) is a map of functors that commutes with χ and δ . We denote by $Aut(F)$

the group of automorphisms of F as a monoidal functor (we will not consider automorphisms of a monoidal functor as a functor).

More generally, if $\mathcal{C}_0$ is a third monoidal category, and $(I, \varepsilon) : \mathcal{C}_0 \rightarrow \mathcal{C}$ is a monoidal functor, we denote by $Aut(F/\mathcal{C}_0)$ (or $Aut(F/I)$) the subgroup of $Aut(F)$ consisting of elements θ such that $\theta_{I(\mathbf{X})}$ is the identity for all objects $\mathbf{X}$ of $\mathcal{C}_0$.

1.3. Closed categories. Let $\mathcal{C}$ be a symmetric monoidal category. If, for any object $\mathbf{X}$ of $\mathcal{C}$, the functor $\mathbf{Y} \mapsto \mathbf{Y} \otimes \mathbf{X}$ has a right adjoint $\mathbf{Z} \mapsto \underline{Hom}(\mathbf{X}, \mathbf{Z})$, then $\mathcal{C}$ is said to be *closed*. Thus, if $\mathcal{C}$ is closed, we have for any three objects $\mathbf{X}$, $\mathbf{Y}$ and $\mathbf{Z}$, $Hom(\mathbf{Y} \otimes \mathbf{X}, \mathbf{Z}) \xrightarrow{\sim} Hom(\mathbf{Y}, \underline{Hom}(\mathbf{X}, \mathbf{Z}))$, functorially in each term. If two maps correspond to each other under this isomorphism, we call each the transpose of the other.

Setting $\mathbf{Y} = \underline{Hom}(\mathbf{X}, \mathbf{Z})$ and applying the isomorphism to the identity map gives rise to the evaluation map $ev = ev_{\mathbf{X}, \mathbf{Z}} : \underline{Hom}(\mathbf{X}, \mathbf{Z}) \otimes \mathbf{X} \rightarrow \mathbf{Z}$. Setting $\mathbf{Y} = \underline{Hom}(\mathbf{W}, \mathbf{Z}) \otimes \underline{Hom}(\mathbf{X}, \mathbf{W})$, and applying the adjunction to the map $ev_{\mathbf{W}, \mathbf{Z}} \circ (id_{\underline{Hom}(\mathbf{W}, \mathbf{Z})} \otimes ev_{\mathbf{X}, \mathbf{W}})$ gives a composition map $\circ_{\mathbf{X}, \mathbf{W}, \mathbf{Z}} : \underline{Hom}(\mathbf{W}, \mathbf{Z}) \otimes \underline{Hom}(\mathbf{X}, \mathbf{W}) \rightarrow \underline{Hom}(\mathbf{X}, \mathbf{Z})$.

Applying the above adjunction with the object $\mathbf{1}$ shows that $\Gamma(\underline{Hom}(\mathbf{X}, \mathbf{Y})) = Hom(\mathbf{X}, \mathbf{Y})$, and that the combined map

$$Hom(\mathbf{Y}, \mathbf{Z}) \times Hom(\mathbf{X}, \mathbf{Y}) = \Gamma(\underline{Hom}(\mathbf{Y}, \mathbf{Z})) \times \Gamma(\underline{Hom}(\mathbf{X}, \mathbf{Y})) \rightarrow \\ \Gamma(\underline{Hom}(\mathbf{Y}, \mathbf{Z}) \otimes \underline{Hom}(\mathbf{X}, \mathbf{Y})) \xrightarrow{\Gamma(\circ)} \Gamma(\underline{Hom}(\mathbf{X}, \mathbf{Z})) = Hom(\mathbf{X}, \mathbf{Z})$$

is the usual composition.

1.4. Example. If $\mathcal{C}$ is a category with finite products, then $\mathcal{C}$ can be viewed as a symmetric monoidal category with the product as $\otimes$, and the usual isomorphisms as constraints. The final object $\mathbf{1}$ with the unique isomorphism $\mathbf{1} \rightarrow \mathbf{1} \times \mathbf{1}$ is the (unique) identity object. We call this the *Cartesian monoidal structure*. A monoidal functor between two such categories is simply a functor that preserves finite products, and a map between two such functors in the monoidal sense is the same as a map in the usual sense. In particular, the global sections functor is a monoidal functor in this case. A category that has all finite (inverse) limits, and is closed with this monoidal structure is called *Cartesian closed* (Mac Lane [17, p. IV.6]).

1.5. Example. If $\mathcal{D}$ is any category, the category $\mathcal{E}(\mathcal{D})$ of functors from $\mathcal{D}$ to itself is a (non-symmetric) monoidal category with $\otimes$ the composition of functors, and ϕ the identity. The monoidal full subcategory of $\mathcal{E}(\mathcal{D})$ consisting of functors that have a right adjoint is closed (with $\underline{Hom}(\mathbf{F}, \mathbf{G}) = \mathbf{R} \circ \mathbf{G}$, where $\mathbf{R}$ is the right adjoint of $\mathbf{F}$). If $\mathcal{D}$ is itself a monoidal category, the assignment $\mathbf{X} \mapsto F_{\mathbf{X}}$, where $F_{\mathbf{X}}(\mathbf{Y}) = \mathbf{X} \otimes \mathbf{Y}$, is a faithful embedding of $\mathcal{D}$ in $\mathcal{E}(\mathcal{D})$, which becomes a monoidal functor with $\phi_{\mathcal{D}}$.

1.6. Example. If R is a commutative ring, the category Mod_R of R -modules, and the subcategory $\mathcal{V}ec_R$ of finitely generated projective R -modules, are symmetric monoidal categories, with the usual tensor product and the usual constraints. Examples of this type will be considered in more details starting from section 3.

1.7. Ind-closed monoidal categories. Let $\mathcal{C}$ be a category, and let $\hat{\mathcal{C}}$ be the category of pre-sheaves (contra-variant functors to the category of sets) on $\mathcal{C}$. Recall that $\text{Ind}(\mathcal{C})$, the category of ind-objects of $\mathcal{C}$, is the full subcategory of $\hat{\mathcal{C}}$ consisting of functors of the form $\varinjlim \mathbf{y}(\mathbf{X}_i)$, where $\mathbf{X}_i$ is a filtering system, and $\mathbf{y} : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ is the Yoneda embedding (cf. Grothendieck et al. [9, p. I] or Artin and Mazur [1] for details on ind- and pro- objects; Gaitsgory and Kazhdan [7] or Kamensky [15] for a short overview). We view $\mathcal{C}$ as a (full) subcategory of $\text{Ind}(\mathcal{C})$. We call a pre-sheaf on $\mathcal{C}$ ind-representable if it is in $\text{Ind}(\mathcal{C})$.

If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, we also denote by $F : \text{Ind}(\mathcal{C}) \rightarrow \text{Ind}(\mathcal{D})$ its extension to the ind categories. In particular, if $\mathcal{C}$ is a monoidal category, we denote by $\otimes$ the extension of the tensor functor to $\text{Ind}(\mathcal{C})$. It is easy to see that the constraints on $\mathcal{C}$ extend to $\text{Ind}(\mathcal{C})$, and so, $\text{Ind}(\mathcal{C})$ is again a (symmetric) monoidal category. If (F, c) is a monoidal functor, it is again easy to see that c extends to give a monoidal structure on (the extended) F . In particular, $\mathcal{C}$ is a monoidal subcategory of $\text{Ind}(\mathcal{C})$.

We say that $\mathcal{C}$ is *ind-closed* if for any two objects $\mathbf{X}$ and $\mathbf{Y}$ of $\mathcal{C}$, the pre-sheaf $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$ given by $\mathbf{Z} \mapsto \text{Hom}(\mathbf{Z} \otimes \mathbf{X}, \mathbf{Y})$ is ind-representable. It follows immediately that $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$ is the right adjoint of $\mathbf{Z} \mapsto \mathbf{Z} \otimes \mathbf{X}$ as a functor on $\text{Ind}(\mathcal{C})$ (however, it is *not* true that $\text{Ind}(\mathcal{C})$ is closed!).

1.8. Proposition. Assume $\mathcal{C}$ has all finite direct limits (or, more generally, that these limits exist in $\text{Pro}(\mathcal{C})$ or in $\text{Ind}(\mathcal{C})$). Then $\mathcal{C}$ is ind-closed if and only if, for any object $\mathbf{X}$ of $\mathcal{C}$, $- \otimes \mathbf{X}$ is right exact (i.e., preserves these limits).

Proof. If $- \otimes \mathbf{X}$ has a right adjoint, then it is right exact. The converse follows from the Grothendieck construction (cf. Mac Lane and Moerdijk [18, p. VII]), which shows that in this situation, $\text{Ind}(\mathcal{C})$ consists precisely of the left exact pre-sheaves (those that take direct limits to the corresponding inverse limits). Since $\mathbf{y}(\mathbf{Y})$ is left exact by definition, $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$ is left exact if $- \otimes \mathbf{X}$ is right exact. $\square$

1.9. Closed monoidal functors. If $\mathcal{C}$ and $\mathcal{D}$ are (ind-) closed, and (F, c) is a monoidal functor, we get, for any two objects $\mathbf{X}, \mathbf{Y}$ of $\mathcal{C}$, a map $F(\underline{\text{ev}}_{\mathbf{X}, \mathbf{Y}}) \circ c_{\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}), \mathbf{X}}$ from $F(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})) \otimes F(\mathbf{X})$ to $F(\mathbf{Y})$, and therefore a map from $F(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))$ to $\underline{\text{Hom}}(F(\mathbf{X}), F(\mathbf{Y}))$. We say that F is *closed* if this map is an isomorphism for any two objects $\mathbf{X}$ and $\mathbf{Y}$ of $\mathcal{C}$.

For example, if R is a commutative ring, and S is a (commutative) R -algebra, then the functor $\mathbf{M} \mapsto S \otimes \mathbf{M}$ from $\mathcal{V}_{ec R}$ to $\mathcal{V}_{ec S}$ (as in example 1.6) is closed. More generally, any monoidal functor between rigid categories (3.1) is closed (cf. Deligne and Milne [5, Prop. 1.9]).

1.10. The rest of this section is devoted to the proof of a result (1.13) that I consider to be an analogue of the existence of definable Galois groups in the context of internal covers (2.5). It will be used in section 2 to recover that result, but is not needed elsewhere in the paper.

Till the end of this section, we will restrict to the case, as in example 1.4, of categories with finite inverse limits, with the Cartesian monoidal structure. Note that if F is a monoidal functor from such a category to a monoidal category $\mathcal{D}$, and F is left exact, then the monoidal structure on $\mathcal{D}$ is also Cartesian, at least when restricted to the essential image of F . We will also assume that monoidal

functors between two such categories preserve all finite inverse limits (rather than just products).

We call a category *ind-Cartesian* if it is ind-closed with this monoidal structure. Thus, any Cartesian-closed category, and in particular any elementary topos, in the sense of Mac Lane and Moerdijk [18], is ind-Cartesian, and any “logical morphism” is a closed monoidal functor. However, the category of definable sets in a theory, considered in section 2, is usually at most ind-Cartesian, and hence is not an elementary topos.

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a left exact functor between two categories. Given an object $\mathbf{T}$ of $\mathcal{D}$, we denote by $\text{End}_{\mathbf{T}}(F)$ the set of natural transformations from $\mathbf{T} \times F$ to F . This is a contra-variant functor in $\mathbf{T}$. We denote by $\mathbf{T} \mapsto \text{Aut}_{\mathbf{T}}(F)$ the subfunctor consisting of those transformations $\mathbf{T} \times F \rightarrow F$ for which the induced map $\mathbf{T} \times F \rightarrow \mathbf{T} \times F$ is an isomorphism. The functors $\mathbf{T} \mapsto \text{End}_{\mathbf{T}}(F)$ and $\mathbf{T} \mapsto \text{Aut}_{\mathbf{T}}(F)$ take values in the categories of monoids and groups, respectively.

As in 1.2, if $I : \mathcal{C}_0 \rightarrow \mathcal{C}$ is another left exact functor, we denote by $\text{End}_{\mathbf{T}}(F/I)$ and by $\text{Aut}_{\mathbf{T}}(F/I)$ the submonoid (subgroup) of $\text{End}_{\mathbf{T}}(F)$ and $\text{Aut}_{\mathbf{T}}(F)$ consisting maps where the induced endomorphism of $\mathbf{T} \times F$ restricts to the identity on $\mathcal{C}_0$.

If $\mathcal{C}$ is a category with finite products, a *pro-group* in $\mathcal{C}$ is a co-filtering system of group objects in $\mathcal{C}$ (viewed as a group object of $\text{Pro}(\mathcal{C})$). We note that an arbitrary group object in $\text{Pro}(\mathcal{C})$ need not be of this form.

1.11. Proposition. Let $\mathcal{C}_0$, $\mathcal{C}$ and $\mathcal{D}$ be categories with all finite inverse limits, and let $I : \mathcal{C}_0 \rightarrow \mathcal{C}$ and $F : \mathcal{C} \rightarrow \mathcal{D}$ be left-exact functors. Assume that $\underline{\text{Hom}}(F(\mathbf{X}), F(\mathbf{Y}))$ exists for all objects $\mathbf{X}$ and $\mathbf{Y}$ of $\mathcal{C}$. Then the functors $\mathbf{T} \mapsto \text{End}_{\mathbf{T}}(F/I)$ and $\mathbf{T} \mapsto \text{Aut}_{\mathbf{T}}(F/I)$ are pro-representable, by a pro-monoid $\mathbf{M}$ and pro-group $\mathbf{G}$, respectively.

In particular, if $H : \mathcal{D} \rightarrow \mathcal{E}$ is any *closed* left-exact functor, then $\Gamma_{\mathcal{E}}(H(\mathbf{M})) = \text{End}(H \circ F/I)$ and $\Gamma_{\mathcal{E}}(H(\mathbf{G})) = \text{Aut}(H \circ F/I)$.

Proof. We first assume that $\mathcal{C}_0$ is empty. Let $\mathcal{J}$ be the following category: An object of $\mathcal{J}$ is a morphism f of $\mathcal{C}$, and a morphism from f to g is a pair of morphisms (u, v) in $\mathcal{C}$ satisfying $v \circ f \circ u = g$. We define a system $\mathbf{M}_i$ in $\mathcal{D}$ indexed by $\mathcal{J}$, as follows: If $f : \mathbf{X} \rightarrow \mathbf{Y}$ is an object of $\mathcal{J}$, $M_f = \underline{\text{Hom}}(F(\mathbf{X}), F(\mathbf{Y}))$. If (u, v) is a morphism from $f : \mathbf{X} \rightarrow \mathbf{Y}$ to $g : \mathbf{Z} \rightarrow \mathbf{W}$, then $M_{(u,v)} : M_f \rightarrow M_g$ is $\underline{\text{Hom}}(\mathbf{Z}, v) \circ \underline{\text{Hom}}(u, \mathbf{Y}) = \underline{\text{Hom}}(u, \mathbf{W}) \circ \underline{\text{Hom}}(\mathbf{X}, v)$. We set $M = \text{Pro}_{\mathcal{J}}(\mathbf{M}_i)$ (the category $\mathcal{J}$ is not co-filtering, but the system can be canonically completed to a co-filtering one, since $\mathcal{D}$ has finite inverse limits).

For $\mathbf{X}$ an object of $\mathcal{C}$, we denote by $\mathbf{X}$ also the object $\text{id}_{\mathbf{X}} : \mathbf{X} \rightarrow \mathbf{X}$ of $\mathcal{J}$. A morphism t from $\mathbf{T}$ to $\mathbf{M}$ corresponds to a suitable collection of maps $t_f : \mathbf{T} \rightarrow \mathbf{M}_f$. We claim that the system $\theta_{\mathbf{X}} : \mathbf{T} \times F(\mathbf{X}) \rightarrow F(\mathbf{X})$ corresponding to the transposed maps of the $t_{\mathbf{X}}$ is a natural map. Indeed, if $g : \mathbf{X} \rightarrow \mathbf{Y}$ is a map in $\mathcal{C}$, then $F(g) \circ \theta_{\mathbf{X}}$ is the image of $t_{\mathbf{X}}$ under the map $(\text{id}_{\mathbf{X}}, g)$ from $\mathbf{X}$ to $g : \mathbf{X} \rightarrow \mathbf{Y}$ in $\mathcal{J}$, while $\theta_{\mathbf{Y}} \circ (1 \times F(g))$ is the image of $t_{\mathbf{Y}}$ under $(g, \text{id}_{\mathbf{Y}})$. Both of these should be equal to t_g , and in particular should be equal to each other.

Conversely, if $\theta : \mathbf{T} \times F \rightarrow F$ is a natural map, we define a map $t : \mathbf{T} \rightarrow \mathbf{M}$ by $t_f = M_{(\text{id}_{\mathbf{X}}, f)} \circ t_{\mathbf{X}}$, where $f : \mathbf{X} \rightarrow \mathbf{Y}$ is an object of $\mathcal{J}$, and $t_{\mathbf{X}}$ is the transpose of $\theta_{\mathbf{X}}$. If (u, v) is a map from $f : \mathbf{X} \rightarrow \mathbf{Y}$ to $g : \mathbf{Z} \rightarrow \mathbf{W}$, then $M_{(u,v)} \circ t_f = M_{(u, v \circ f)} \circ t_{\mathbf{X}}$. This map corresponds under transposition to the map $\mathbf{T} \times F(\mathbf{Z}) \rightarrow F(\mathbf{W})$ given by $(\text{id}_{\mathbf{T}} \times (v \circ f)) \circ \theta_{\mathbf{X}} \circ (\text{id}_{\mathbf{T}} \times F(u))$. Since θ is natural, $\theta_{\mathbf{X}} \circ (\text{id}_{\mathbf{T}} \times F(u)) = F(u) \circ \theta_{\mathbf{Z}}$,

and substituting this in the last equality, we get $(id_{\mathbf{T}} \times (v \circ f \circ u)) \circ \theta_{\mathbf{Z}} = (id_{\mathbf{T}} \times g) \circ \theta_{\mathbf{Z}}$, which corresponds to $M_{(id_{\mathbf{Z}}, g)} \circ t_{\mathbf{Z}} = t_g$.

This concludes the proof that $\mathbf{M}$ represents $End_{\mathbf{T}}(F)$. Since this functor is a monoid, so is $\mathbf{M}$. If $I : \mathcal{C}_0 \rightarrow \mathcal{C}$ is now as in the statement, and M_0 is the pro-object corresponding to $F \circ I$, then we have an obvious map (“restriction”) from M to M_0 , and the monoid that represents $End_{\mathbf{T}}(F/I)$ is the pullback of this map under the map $\mathbf{1} \rightarrow \mathbf{M}_0$ corresponding to the monoid identity. Likewise, $\mathbf{G}$ is the sub-object of $\mathbf{M}$ of “invertible elements”, again given by a suitable pullback.

The statement that $\mathbf{M}$ and $\mathbf{G}$ are a pro-monoid and a pro-group follow since one could present $\mathcal{C}$ as a union of “finitely generated” sub-categories, so that on each step the system is equivalent to a finite one.

The last statement follows since $H(\mathbf{M}_f)$ is identified with the corresponding construction for the functor $H \circ F$ when H is closed (and by applying the statement with $\mathbf{T} = \mathbf{1}$). $\square$

1.12. Internal covers. In model theory, one is interested in the automorphisms of functors of the form $\Gamma \circ F$ (for particular functors F). Proposition 1.11 is not very helpful in this situation, since Γ is almost never closed. However, a similar conclusion holds for Γ under the assumption of internality.

By analogy with the model-theoretic terminology, we say that a pair of left-exact functors $I : \mathcal{C}_0 \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}_0$ between ind-closed categories is an *internal cover* (or that $\mathcal{C}$ is an internal cover of $\mathcal{C}_0$) if they are closed, and $F \circ I$ is isomorphic to the identity.

We denote by $G = Aut(\mathcal{C}/\mathcal{C}_0)$ the pro-group provided by 1.11, with F the embedding of $\mathcal{C}$ in $Ind(\mathcal{C})$. Thus, G is a projective system of groups in $Ind(\mathcal{C})$.

1.13. Theorem. Let $I : \mathcal{C}_0 \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}_0$ be an internal cover. Then for any left-exact $H : \mathcal{C}_0 \rightarrow \mathcal{D}$, the map

$$Aut(H \circ F/I) \rightarrow Aut(\Gamma_{\mathcal{D}} \circ H \circ F/I)$$

(obtained by applying $\Gamma_{\mathcal{D}}$) is an isomorphism. Hence, there is a pro-group $\mathbf{G}$ in $Ind(\mathcal{C})$, such that

$$\Gamma_{\mathcal{D}}(H(F(\mathbf{G}))) = Aut(\Gamma_{\mathcal{D}} \circ H \circ F/I)$$

for any closed H .

Proof. The proof closely follows the proof of the analogous result 2.5 in model theory. We identify $F \circ I$ with the identity, by fixing an isomorphism, and set $R = \Gamma_{\mathcal{D}} \circ H \circ F$. Let θ be an automorphism over I of R , let $\mathbf{Q}$ be an object of $\mathcal{C}$, and set $\mathbf{X} = \underline{Hom}(\mathbf{Q}, I(F(\mathbf{Q})))$. We will prove that $\theta_{\mathbf{Q}}$ is represented by a (unique) element of $R(\mathbf{X})$.

We first note that

$$R(\mathbf{X}) = \Gamma_{\mathcal{D}}(H(\underline{Hom}(F(\mathbf{Q}), F(I(F(\mathbf{Q})))))) = \Gamma_{\mathcal{D}}(H(\underline{Hom}(F(\mathbf{Q}), F(\mathbf{Q}))))$$

since F is closed, and $F \circ I$ is the identity. $R(\mathbf{X})$ therefore contains an element 1, corresponding to the identity map on $F(\mathbf{Q})$. Also, the evaluation map

$$ev : R(\mathbf{X}) \times R(\mathbf{Q}) \rightarrow R(I(F(\mathbf{Q}))) = R(\mathbf{Q})$$

is preserved by θ , since it comes from $\mathcal{C}$. Finally, θ is constant on the co-domain of ev , since it has the form $R(I(\mathbf{C}))$ (for $\mathbf{C} = F(\mathbf{Q})$). Hence, if $f \in R(\mathbf{X})$ and $u \in R(\mathbf{Q})$, then $\theta_{\mathbf{X}}(f)(\theta_{\mathbf{Q}}(u)) = f(u)$ (where $f(u) = ev(f, u)$). In particular, for

$f = 1$ we get that $\theta_{\mathbf{X}}(1)$ is invertible, and $\theta_{\mathbf{Q}} = (\theta_{\mathbf{X}}(1))^{-1}$. In other words, $\theta_{\mathbf{Q}}^{-1}$ is given by $H(F(\text{ev}_{\mathbf{Q}, I(F(\mathbf{Q}))})) \circ (\theta_{\mathbf{X}}(1) \times \text{Id})$, a morphism in $\mathcal{D}$. $\square$

In the model theoretic setting, the group $\text{Aut}(\mathcal{C}/\mathcal{C}_0)$ above is actually a pro-group in $\mathcal{C}$, rather than $\text{Ind}(\mathcal{C})$. This seems to require additional assumptions.

1.14. We mention two alternative formulations of 1.13.

1.14.1. Assume we are given functors $I_1 : \mathcal{C}_0 \rightarrow \mathcal{C}_1$, $J : \mathcal{C}_1 \rightarrow \mathcal{C}$ and $F : \mathcal{C} \rightarrow \mathcal{C}_1$ such that J and F form an internal cover, and assume that $I = J \circ I_1$ is closed. Then the theorem holds for I and F (for any left-exact $H : \mathcal{C}_1 \rightarrow \mathcal{D}$), since $\text{Aut}(\mathbf{T}/I) = \text{Aut}(\mathbf{T}/J)$ for any functor $\mathbf{T}$.

In particular, if $I : \mathcal{C}_0 \rightarrow \mathcal{C}$ is closed, we can take $\mathcal{C}_1$ to be the essential image of I , and J the inclusion. In this case, F is a functor from $\mathcal{C}$ to itself. We will call such a pair an internal cover as well.

1.14.2. If $\mathcal{C}$ is an ind-closed category, and $F' : \mathcal{C} \rightarrow \mathcal{C}$ is a closed functor such that $F' \circ F'$ is isomorphic to F' , we may apply the previous part with $F = I = F'$. Conversely, in the situation of 1.13, $F' = I \circ F$ is closed and isomorphic to $F' \circ F'$.

2. INTERNALITY IN MODEL THEORY

In this section, we recall some notions from model theory. Our goal is to explain the notions of internality and the associated Galois group. We then demonstrate how part of this result can be interpreted in the language of the previous section. We shall, for the most part, follow the presentation and terminology from Hrushovski [12].

2.1. We assume standard notions from logic to be known. We briefly recall that a *theory* is collection of statements (axioms) written in a fixed formal language, and a *model* of the theory $\mathcal{T}$ is a structure consisting of an interpretation of the symbols in the language of $\mathcal{T}$, in which all the axioms of $\mathcal{T}$ hold. For example, the theory ACF of algebraically closed fields can be written in a language containing symbols $(0, 1, +, -, \cdot)$, and a model of this theory is an algebraically closed field.

A *formula* is written in the same formal language, but has *free variables*, into which elements of the model can be plugged. For the example of ACF above, any finite collection of polynomial equations and inequalities can be viewed as a formula, but there are other formulas, involving quantifiers. Any such formula $\phi(x_1, \dots, x_n)$ (where $x_1, \dots, x_n$ contain all free variables of ϕ) thus determines a subset $\phi(M)$ of M^n , for any model M , namely, the set of all tuples $\bar{a}$ for which $\phi(\bar{a})$ holds. Two formulas ϕ and ψ are *equivalent* if $\phi(M) = \psi(M)$ for all models M . An equivalence class under this relation is called a *definable set*.

If M is the model of some theory, the set of all statements (in the underlying language) that are true in M is a theory $\mathcal{T}(M)$, and M is a model of $\mathcal{T}(M)$.

We will assume all our theories to be multi-sorted, i.e., the variables of a formula can take values in any number of disjoint sets. In fact, if $\mathbf{X}$ and $\mathbf{Y}$ are sorts, we view $\mathbf{X} \times \mathbf{Y}$ as a new sort. In particular, any definable set is a subset of some sort. By a statement such as $a \in M$ we will mean that a is an element of one of the sorts.

2.2. Definable closure and automorphisms. A *definable function* f from a definable set $\mathbf{X}$ to another definable set $\mathbf{Y}$ is a definable subset of $\mathbf{X} \times \mathbf{Y}$ that determines the graph of a function $f_M : \mathbf{X}(M) \rightarrow \mathbf{Y}(M)$ for each model M . If M is a model, $A \subseteq \mathbf{X}(M)$ any subset, and $b \in \mathbf{Y}(M)$, then b is *definable* over A if there is a definable function $f : \mathbf{X}^n \rightarrow \mathbf{Y}$ for some n , such that $f(\bar{a}) = b$ for some $\bar{a} \in A^n$ (note that even though we are working in M , it is $\mathcal{T}$ that should think that f is a function, rather than just $\mathcal{T}(M)$). A subset is *definably closed* if it is closed under definable functions. The *definable closure* $\text{dcl}(A)$ of a subset $A \subseteq M$ is the smallest definably closed subset of M containing A . We denote by $\mathbf{Y}(A)$, the set of all elements in $\mathbf{Y}(M)$ definable over A .

We note that if $\mathcal{T}$ is complete, then the definable closure $\mathbf{X}(0)$ of the empty set does not depend on the model. For general $\mathcal{T}$, we denote by $\mathbf{X}(0)$ the set of definable subsets of $\mathbf{X}$ containing one element.

More generally, a formula over A is a formula $\phi(x, a)$, where $\phi(x, y) \subseteq \mathbf{X} \times \mathbf{Y}$ is a regular formula, and $a \in \mathbf{Y}(A)$. It defines a subset of $\mathbf{X}(M)$ for any model M containing A , in the same way as regular formulas do.

An automorphism of a model M is a bijection φ from M to itself such that the induced bijection $\varphi^n : M^n \rightarrow M^n$ preserves any definable subset $\mathbf{X}(M) \subseteq M^n$. If $A \subseteq M$ is a set of parameters, the automorphism φ is *over* A if it fixes all elements of A . The group of all automorphisms of M over A is denoted by $\text{Aut}(M/A)$. It is clear that $\mathbf{X}(A)$ is fixed pointwise by any automorphism over A . M is called *homogeneous* if, conversely, for any $A \subseteq M$ of cardinality smaller than the cardinality of M , for any definable set $\mathbf{X}$, any element of $\mathbf{X}(M)$ fixed by all of $\text{Aut}(M/A)$ is in $\mathbf{X}(A)$ (computed in $\mathcal{T}(M)$). Homogeneous models are known to exist (cf. Sacks [29, Ch. 20]).

2.3. Imaginaries and interpretations. A *definable equivalence relation* on a definable set $\mathbf{X}$ is a definable subset of $\mathbf{X} \times \mathbf{X}$ that determines an equivalence relation in any model. The theory *eliminates imaginaries* if any equivalence relation has a quotient, i.e., any equivalence relation can be represented as $f(x) = f(y)$ for some definable function f on $\mathbf{X}$.

A *definition* of a theory $\mathcal{T}_1$ in another theory $\mathcal{T}_2$ is specified by giving, for each sort $\mathbf{X}$ of $\mathcal{T}_1$ a definable set $\mathbf{X}_2$ of $\mathcal{T}_2$, and for each atomic relation $\mathbf{Y} \subseteq \mathbf{X}$ of $\mathcal{T}_1$ a definable subset $\mathbf{Y}_2$ of $\mathbf{X}_2$, such that for any model M_2 of $\mathcal{T}_2$, the sets $\mathbf{X}_2(M_2)$ form a model M_1 of $\mathcal{T}_1$ (when interpreted as a $\mathcal{T}_1$ structure in the obvious way). It follows that any definable set of $\mathcal{T}_1$ determines a definable set in $\mathcal{T}_2$, and M_1 can be viewed as a subset of M_2 .

For any theory $\mathcal{T}$ there is a theory $\mathcal{T}^{eq}$ and a definition of $\mathcal{T}$ in $\mathcal{T}^{eq}$, where $\mathcal{T}^{eq}$ eliminates imaginaries, and the definition is universal with this property. Any model of $\mathcal{T}$ can be expanded uniquely to a model of $\mathcal{T}^{eq}$. Replacing $\mathcal{T}$ with $\mathcal{T}^{eq}$, if necessary, we shall usually assume that *all our theories eliminate imaginaries*.

An *interpretation* of $\mathcal{T}_1$ in $\mathcal{T}_2$ is a definition of $\mathcal{T}_1$ in $\mathcal{T}_2^{eq}$. Hence, with the above convention, interpretations and definitions coincide. Given an interpretation of $\mathcal{T}_1$ in $\mathcal{T}_2$, we view the definable sets of $\mathcal{T}_1$ as definable sets in $\mathcal{T}_2$ via the interpretation.

If M is a model of a theory $\mathcal{T}$, and $A \subseteq M$, we denote by $\mathcal{T}_A$ the theory obtained by adding constants for A to the language, and the axioms satisfied for A in M (in particular, $\mathcal{T}_A$ is complete). There is an obvious interpretation of $\mathcal{T}$ in $\mathcal{T}_A$. The model M is in a natural way a model of $\mathcal{T}_A$, and for any definable set $\mathbf{X}$ of $\mathcal{T}$, $\mathbf{X}(A)$ in the sense of $\mathcal{T}$ is identified with $\mathbf{X}(0)$ in $\mathcal{T}_A$.

2.4. Internal covers. An interpretation of $\mathcal{T}_0$ in $\mathcal{T}$ is *stably embedded* if any subset of the sorts of $\mathcal{T}_0$ definable in $\mathcal{T}$ with parameters from $\mathcal{T}$ is also definable in $\mathcal{T}_0$ with parameters from $\mathcal{T}_0$.

A stably embedded interpretation of $\mathcal{T}_0$ in $\mathcal{T}$ is an *internal cover* (of $\mathcal{T}_0$) if any large enough model M of $\mathcal{T}$ has a small subset $A \subseteq M$ such that $\text{dcl}(M_0 \cup A) = M$. $\mathcal{T}$ itself is also called an *internal cover* of $\mathcal{T}_0$. A set A as above will be called a set of *internality parameters*, and will always be taken to be definably closed. We denote by A_0 the restriction of A to $\mathcal{T}_0$. (In the language of Hrushovski [12], this implies that the corresponding groupoid in $\mathcal{T}_0$ is equivalent to a groupoid with objects over A_0 .)

We note that by compactness, a finite number of elements of A suffices to define all elements of a given sort of $\mathcal{T}$ over M_0 . Thus, if $\mathcal{T}$ has a finite number of sorts that do not come from $\mathcal{T}_0$ over which everything is definable, then A can be taken to be generated by one tuple a . In general, A can be thought of as an element of a pro-definable set.

2.5. Theorem. (Hrushovski [12, prop. 1.5] and Hrushovski [11, appendix B]) Let $\mathcal{T}$ be an internal cover of $\mathcal{T}_0$. There is a pro-group $\mathbf{G}$ in $\mathcal{T}$, together with a definable action $m_Q : \mathbf{G} \times \mathbf{Q} \rightarrow \mathbf{Q}$ of $\mathbf{G}$ on every definable set $\mathbf{Q}$ of $\mathcal{T}$, such that for any model M of $\mathcal{T}$, $\mathbf{G}(M)$ is identified with $\text{Aut}(M/M_0)$ through this action (with M_0 the restriction of M to $\mathcal{T}_0$).

Furthermore, given a set of internality parameters A (whose restriction to $\mathcal{T}_0$ is A_0), there is a Galois correspondence between A -definable pro-subgroups of $\mathbf{G}$ and definably closed subsets $A_0 \subseteq B \subseteq A$. Such a subset B_H for a subgroup $\mathbf{H}$ is always of the form $\mathbf{C}(A)$, where $\mathbf{C}$ is an A -definable ind-set in $\mathcal{T}$, and $\mathbf{H}$ is the subgroup of $\mathbf{G}$ fixing $\mathbf{C}(M)$ pointwise. If $\mathbf{H}$ is normal, then $\mathbf{G}/\mathbf{H}(M)$ is identified with $\text{Aut}(\mathbf{C}(M)/M_0)$.

Here, a *pro-group* is a filtering inverse system of definable groups.

2.6. In fact, the result is slightly stronger. With notation as above, the assumption that $\mathcal{T}_0$ is stably embedded implies that any automorphism of M_0 fixing A_0 can be extended (uniquely) to an automorphism of M fixing A . In other words, we have a split exact sequence

$$0 \rightarrow \mathbf{G}(M) \rightarrow \text{Aut}(M/A_0) \rightarrow \text{Aut}(M_0/A_0) \rightarrow 0 \quad (1)$$

where $\mathbf{G}$ is as provided by 2.5.

For any definably closed subset $B_0 \subseteq M_0$ containing A_0 , let $B = \text{dcl}(A \cup B_0)$. We claim that $\mathbf{G}(B) = \text{Aut}(B/B_0)$. Indeed, any $g \in \mathbf{G}(B)$ acts as an automorphism, preserves B as a set (since B is definably closed), and fixes B_0 pointwise, so $\mathbf{G}(B) \subseteq \text{Aut}(B/B_0)$. Conversely, since B_0 is definably closed and contains A_0 , $B \cap M = B_0$, and so any automorphism of B over B_0 extends to an automorphism of B over M_0 . Since M_0 is stably embedded, this automorphism extends to an automorphism of M . Thus, any element of $\text{Aut}(B/B_0)$ is represented by some $g \in \mathbf{G}(M)$. Since it is fixed by any automorphism fixing B pointwise, it is in fact in $\mathbf{G}(B)$.

2.7. We make the assumption that a set A as above can be found (in some model) such that the corresponding set A_0 is (the definable closure of) the empty set. This is not a real assumption in the current context, since the results will hold in general for the theory with parameters from A .

Likewise, the interpretation of $\mathcal{T}_0$ in $\mathcal{T}$ factors through a maximal extension $\mathcal{T}_1$ of $\mathcal{T}_0$ (i.e., $\mathcal{T}_1$ is the theory, in $\mathcal{T}$, of the definable sets coming from $\mathcal{T}_0$). We assume from now on that $\mathcal{T}_1 = \mathcal{T}_0$.

Combining the two assumptions, we get that $B_0 \mapsto B = \text{dcl}(A \cup B_0)$ gives an equivalence between definably closed sets B_0 of $\mathcal{T}_0$ and definably closed sets B containing A of $\mathcal{T}$. In particular, we have a definable group $\mathbf{G}_A$ in $\mathcal{T}_0$. The following proposition says that all definable group action of $\mathbf{G}_A$ in $\mathcal{T}_0$ come from the canonical action of $\mathbf{G}$ in $\mathcal{T}$.

2.8. Proposition. Assume 2.7, and let $a : \mathbf{G}_A \times \mathbf{D} \rightarrow \mathbf{D}$ be a definable group action in $\mathcal{T}_0$. There is a definable set $\mathbf{X}_{\mathbf{D}}$ in $\mathcal{T}$, and an A -definable isomorphism of $\mathbf{G}_A$ actions from $\mathbf{X}_{\mathbf{D}}$ to $\mathbf{D}$. If $\mathbf{D}$ and $\mathbf{E}$ are two such $\mathbf{G}_A$ -sets, and $f : \mathbf{D} \rightarrow \mathbf{E}$ is a definable map of $\mathbf{G}_A$ sets in $\mathcal{T}_0$, it also comes from a definable map $F : \mathbf{X}_{\mathbf{D}} \rightarrow \mathbf{X}_{\mathbf{E}}$.

Proof. The proof of 2.5 produces a definable $\mathbf{G}$ -torsor $\mathbf{X}$ in $\mathcal{T}$, which is 0-definable with our assumptions. There is an A -definable isomorphism $f_b : \mathbf{G} \rightarrow \mathbf{G}_A$, which, composed with a , defines an A -definable action $c_b : \mathbf{G} \times \mathbf{D} \rightarrow \mathbf{D}$. We set $\mathbf{X}_{\mathbf{D}} = \mathbf{X} \times_{\mathbf{G}} \mathbf{D}$, i.e., $\mathbf{X}_{\mathbf{D}}$ is the set of pairs $(x, d) \in \mathbf{X} \times \mathbf{D}$, up to the equivalence $(gx, d) = (x, c_b(g, d))$ for $g \in \mathbf{G}$. Since $\mathbf{G}_A$ is in $\mathcal{T}_0$, $f_{hb}(hgh^{-1}) = f_b(g)$ for all $h, g \in G$, so the equivalence relation is invariant under the action of G , and hence $\mathbf{X}_{\mathbf{D}}$ is definable without parameters.

If $b \in \mathbf{X}(A)$ is any element (known to exist), the map $[b, d] \mapsto d$ is an isomorphism of $\mathbf{G}_A$ actions from $\mathbf{X}_{\mathbf{D}}$ to $\mathbf{D}$. The proof for maps is similar. $\square$

2.9. We now give a partial proof of 2.5, using 1.13, partial in that it produces only an pro-ind-definable group, rather than an pro-group. The rest of this section will not be used later.

To each theory $\mathcal{T}$ we associate the category $D_{\mathcal{T}}$ of definable sets and definable maps between them. This category is closed under finite inverse limits and under co-products (this requires that $\mathcal{T}$ has a least two constants, which we assume, and which is automatic if $\mathcal{T}$ eliminates imaginaries). We write $\mathbf{1}$ for the one-element set, and $\mathbf{2}$ for its co-product with itself.

We list the categorical interpretation of some of the notions mentioned above. Most of this can be found in much greater detail (and in a slightly different language) in Makkai and Reyes [19]. Let $\mathcal{T}$ be a theory, not necessarily with elimination of imaginaries.

2.9.1. $\mathcal{T}$ eliminates imaginaries if and only if $D_{\mathcal{T}}$ is ind-Cartesian (in the sense of 1.10), if and only if the co-equaliser of any two definable maps exists in $\text{Pro}(D_{\mathcal{T}})$.

Proof. Assume $D_{\mathcal{T}}$ is ind-Cartesian, and let $\mathbf{E} \subseteq \mathbf{X} \times \mathbf{X}$ be an equivalence relation. The corresponding characteristic function $\chi_{\mathbf{E}} : \mathbf{X} \times \mathbf{X} \rightarrow \mathbf{2}$ corresponds, by assumption, to some $t : \mathbf{X} \rightarrow \mathcal{P}(\mathbf{X}) := \underline{\text{Hom}}(\mathbf{X}, \mathbf{2})$. If $\mathcal{P}(\mathbf{X})$ is represented by some system $(\mathbf{Y}_a, f_{a,b})$, the map t is represented by some $t_a : \mathbf{X} \rightarrow \mathbf{Y}_a$, and so the relation $\mathbf{E}(x, y)$ is equivalent to the intersection over all b of the conditions $f_{a,b}(t_a(x)) = f_{a,b}(t_a(y))$, hence, by compactness, by one such condition. The map $f_{a,b} \circ t_a$ onto its image is then the quotient.

If $\mathcal{T}$ eliminates imaginaries, and $f, g : \mathbf{X} \rightarrow \mathbf{Y}$ are two definable maps, let I be the collection of definable equivalence relations $\mathbf{E}$ on $\mathbf{Y}$ such that $\forall x(\mathbf{E}(f(x), g(x)))$ holds. Since I is non-empty and closed under intersections, we get an inverse

system $\mathbf{Z} = \text{Pro}_{\mathbf{E} \in I}(\mathbf{Y}/\mathbf{E})$, together with a map $h : \mathbf{Y} \rightarrow \mathbf{Z}$, which is clearly the co-equalizer.

Finally, if $\text{Pro}(D_{\mathcal{T}})$ has all co-equalisers, then it has all finite co-limits, and hence is ind-Cartesian by 1.8 (Alternatively, the Hom sets can be constructed explicitly under the assumption of EI). $\square$

2.9.2. A model M of $\mathcal{T}$ determines a functor $\mathbf{X} \mapsto \mathbf{X}(M)$ from $D_{\mathcal{T}}$ to the category of sets, which preserves all inverse limits, co-products, and co-equalisers of equivalence relations (these are called “logical functors” in Makkai and Reyes [19]). Conversely, any such functor is isomorphic to a model. An elementary map from M to N as models is the same as a map of the corresponding functors. A subset A of M is definably closed if and only if it defines a left-exact sub-functor of M that preserves co-products.

2.9.3. Similarly, a definition of the theory $\mathcal{T}_1$ in another theory $\mathcal{T}_2$ is the same as a logical functor I (in the same sense as above) from $D_{\mathcal{T}_1}$ to $D_{\mathcal{T}_2}$. Given a model M of $\mathcal{T}_2$ (viewed as a functor), the corresponding model of $\mathcal{T}_1$ is $M \circ I$.

2.9.4. Let $I : D_{\mathcal{T}_1} \rightarrow D_{\mathcal{T}_2}$ be a definition, where $\mathcal{T}_1$ eliminates imaginaries, and let I^* be the induced functor from models of $\mathcal{T}_2$ to models of $\mathcal{T}_1$. Then I is an equivalence of categories if and only if I^* is.

Proof. Assume that I^* is an equivalence, and let $\mathbf{X}$ be a definable set of $\mathcal{T}_2$. If $\mathbf{X} \subseteq I(\mathbf{Y})$ for some $\mathbf{Y}$, then for any two models M and N of $\mathcal{T}_2$ such that $M \circ I = N \circ I$, we have $M(\mathbf{X}) = N(\mathbf{Y})$ (by equivalence). Hence by Beth’s definability theorem, $\mathbf{X} = I(\mathbf{X}_0)$ for some $\mathbf{X}_0$.

If $\mathbf{X}$ is not contained in any $I(\mathbf{Y})$, but there is a surjective map $f : I(\mathbf{Y}) \rightarrow \mathbf{X}$, the kernel is a definable subset of $I(\mathbf{Y})^2$, hence by the previous case comes from $\mathcal{T}_1$. Since $\mathcal{T}_1$ eliminates imaginaries, the quotient $\mathbf{X}$ also comes from $\mathcal{T}_1$.

If there is no surjective map as above, then a model of $\mathcal{T}_2$ has automorphisms over the restriction to $\mathcal{T}_1$, contradicting the equivalence. This shows that I is essentially surjective. It is full since any map is a definable set, and it is faithful since for equality of definable sets is first order. This shows that I is an equivalence. The other direction is trivial. $\square$

2.9.5. If A is a definably closed subset of a model M of the theory $\mathcal{T}$, we denote by I_A the functor from $D_{\mathcal{T}}$ to $D_{\mathcal{T}_A}$ corresponding to the natural interpretation.

Assume $\mathcal{T}$ eliminates imaginaries. Then for any such A , the functor I_A is closed (in the sense of 1.9).

Proof. A definable map f in $\mathcal{T}_A$ from $\mathbf{Z}$ to $I_A(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))$ is given by formulas $\phi(w, z) \subseteq \mathbf{W} \times \mathbf{Z}$ and $\mathbf{F} \subseteq \mathbf{W} \times \mathbf{Z} \times I_A(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))$, such that $\mathbf{Z} = \phi(a, -)$, $f = \mathbf{F}(a, -, -)$, $a \in \mathbf{W}(A)$ and $\mathbf{F}, \mathbf{W}, \phi$ are in $\mathcal{T}$. There is a $\mathcal{T}$ definable subset $\mathbf{W}_0 \subseteq \mathbf{W}$ such that $\mathbf{F}(w, -, -)$ is a function from $\phi(w, -)$ to $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$ for all $w \in \mathbf{W}_0$. We may assume that $\mathbf{W} = \mathbf{W}_0$, so that $\mathbf{F}$ gives a $\mathcal{T}$ definable function from ϕ to $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$. Thus it corresponds to a map $\tilde{F} : \phi \times \mathbf{X} \rightarrow \mathbf{Y}$. The restriction $\tilde{f} : \mathbf{Z} \times \mathbf{X} \rightarrow \mathbf{Y}$ of this map to $\phi(a, -) \times \mathbf{X}$ corresponds to the original map f .

Reversing this argument gives, for each map $\tilde{f} : \mathbf{Z} \times \mathbf{X} \rightarrow \mathbf{Y}$ in $\mathcal{T}_A$ a corresponding map $f : \mathbf{Z} \rightarrow I_A(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))$. This shows that $I_A(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))$ represents the correct functor in $\mathcal{T}_A$, and therefore coincides with $\underline{\text{Hom}}(I_A(\mathbf{X}), I_A(\mathbf{Y}))$.

(More abstractly, $\mathcal{T}_A$ can be viewed as a full sub-category of $\text{Pro}(\mathcal{T})$, and I_A as the inclusion of $\mathcal{T}$ in $\text{Pro}(\mathcal{T})$. This inclusion is always closed.) $\square$

2.9.6. The functor $\mathbf{X} \mapsto \mathbf{X}(0)$ on the definable subsets of a theory $\mathcal{T}$ coincides with $\mathbf{X} \mapsto \Gamma(\mathbf{X}) = \text{Hom}(1, \mathbf{X})$. If A is a definably closed subset of a model M , then it coincides with $\mathbf{X} \mapsto \Gamma_A(I_A(\mathbf{X}))$ as a functor (where Γ_A is global sections in $\mathcal{T}_A$).

2.9.7. Assume that $\mathcal{T}$ eliminates imaginaries. Then for any model M and any definable sets $\mathbf{X}$ and $\mathbf{Y}$ of $\mathcal{T}$, $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})(M)$ is the set of M -definable maps from $\mathbf{X}$ to $\mathbf{Y}$. More generally, this holds when M is a definably closed subset of a model. In particular, $\mathcal{P}(\mathbf{X})(A)$ is the set of A -definable subsets of $\mathbf{X}$.

Proof. According to 2.9.6 and 2.9.5,

$$\begin{aligned} \underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})(M) &= \Gamma_M(I_M(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))) = \\ &= \text{Hom}_M(1, \underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})) = \text{Hom}_M(\mathbf{X}, \mathbf{Y}) \end{aligned} \quad (2)$$

$\square$

2.9.8. Let $I : D_{\mathcal{T}_1} \rightarrow D_{\mathcal{T}_2}$ be the exact functor corresponding to an interpretation, where both theories eliminate imaginaries. Then the interpretation is stably embedded if and only if I is closed.

Proof. Assume that the interpretation is stably embedded, let $\mathbf{X}, \mathbf{Y}$ be definable sets in $\mathcal{T}_1$, and let M be a model of $\mathcal{T}_2$. The map $t : I(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})) \rightarrow \underline{\text{Hom}}(I(\mathbf{X}), I(\mathbf{Y}))$ gives a map

$$t : I(\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y}))(M) \rightarrow \underline{\text{Hom}}(I(\mathbf{X}), I(\mathbf{Y}))(M)$$

By 2.9.7, the co-domain is the set of M -definable maps from $I(\mathbf{X})$ to $I(\mathbf{Y})$ in $\mathcal{T}_2$, while the domain is the set of definable maps from $\mathbf{X}$ to $\mathbf{Y}$ in $\mathcal{T}_1$ with parameters from the model $M \circ I$ (and t assigns to each map the same map viewed as a map between $I(\mathbf{X})$ and $I(\mathbf{Y})$). Since $\mathcal{T}_1$ asserts that all maps in $\underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$ are distinct, and so the same has to hold in $\mathcal{T}_2$. Therefore, t is injective. Since $\mathcal{T}_1$ is stably embedded, any M -definable map between $I(\mathbf{X})$ and $I(\mathbf{Y})$ is definable with parameters in $M \circ I$, hence t is surjective. Since t is a bijection in every model, θ is an isomorphism.

Assume conversely that I is closed, let $\mathbf{X}$ be a definable set in $\mathcal{T}_1$, and let M be a model of $\mathcal{T}_2$. Any M -definable subset $\mathbf{Z} \subseteq I(\mathbf{X})$ (in $\mathcal{T}_2$), corresponds to an M -point of $\mathcal{P}(I(\mathbf{X}))$ (by 2.9.7), hence to an $M \circ I$ point of $\mathcal{P}(\mathbf{X})$, which can be used to define $\mathbf{Z}$. $\square$

2.10. **Proof of 2.5.** Let $I : \mathcal{T}_0 \rightarrow \mathcal{T}$ be an internal cover. By assumption, it is stably embedded, and therefore closed by 2.9.8.

Let A be as in 2.7. As explained there, definable closure gives an equivalence between models of $\mathcal{T}$ containing A , and models of $\mathcal{T}_0$. The former are precisely models of $\mathcal{T}_A$. Since $I_A \circ I : \mathcal{T}_0 \rightarrow \mathcal{T}_A$ (with I_A as in 2.9.5) induces an equivalence on the categories of models, it is itself an equivalence (2.9.4). Let $F = T \circ I_A$, where T is an inverse of $I_A \circ I$. Since F is trivially closed, the pair (I, F) forms an internal cover in the sense of 1.14.1.

Hence by 1.13, there is a pro-group $\mathbf{G}$ in $\text{Ind}(\mathcal{T})$ such that $\Gamma_{\mathcal{D}}(H(F(\mathbf{G}))) = \text{Aut}(\Gamma_{\mathcal{D}} \circ H \circ F/I)$. Setting $\mathcal{D} = \mathcal{T}_{0M}$, and $H = I_M : \mathcal{T}_0 \rightarrow \mathcal{T}_{1M}$ (for any model M

of $\mathcal{T}_0$), and using 2.9.6, we get $G(M \circ F) = \text{Aut}(M \circ F/M)$. Since any model of $\mathcal{T}$ has this form, the theorem is proved.

2.11. Remark. One advantage of this formulation is that the category of definable sets in $\mathcal{T}$ can be replaced by another category. For example, taking particular sub-categories of this category, one easily recovers most of Kamensky [14, thm. 19].

3. TANNAKIAN CATEGORIES

In this section we review, without proofs, the definitions and basic properties of Tannakian categories. The proofs, as well as more details (originally from Saavedra-Rivano [28]) can be found in the first two sections of Deligne and Milne [5]. The section contains no new results (though the terminology is slightly different).

3.1. Definition. Let $\mathcal{C}$ be a closed symmetric monoidal category, and let $\mathbf{1}$ be a fixed identity object. The *dual* $\check{\mathbf{X}}$ of an object $\mathbf{X}$, is the object $\underline{\text{Hom}}(\mathbf{X}, \mathbf{1})$. The adjunction map together the commutativity constraint give rise to a map $\mathbf{X} \rightarrow \check{\check{\mathbf{X}}}$. $\mathbf{X}$ is called *reflexive* if this map is an isomorphism.

For any objects $\mathbf{X}_1, \mathbf{X}_2$ and $\mathbf{Y}_1, \mathbf{Y}_2$, the adjunction also gives maps

$$\underline{\text{Hom}}(\mathbf{X}_1, \mathbf{Y}_1) \otimes \underline{\text{Hom}}(\mathbf{X}_2, \mathbf{Y}_2) \rightarrow \underline{\text{Hom}}(\mathbf{X}_1 \otimes \mathbf{X}_2, \mathbf{Y}_1 \otimes \mathbf{Y}_2)$$

$\mathcal{C}$ is said to be *rigid* if all objects are reflexive, and the above map is always an isomorphism.

We note that setting $\mathbf{Y}_1 = \mathbf{X}_2 = \mathbf{1}$ in the above map, we get an isomorphism $\check{\mathbf{X}} \otimes \mathbf{Y} \rightarrow \underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$. In particular, a map $f : \mathbf{X} \rightarrow \mathbf{Y}$ corresponds to a global section $\mathbf{1} \rightarrow \underline{\text{Hom}}(\mathbf{X}, \mathbf{Y})$, hence to a map $\check{f} : \mathbf{1} \rightarrow \check{\mathbf{X}} \otimes \mathbf{Y}$.

3.2. The rank of an object. Let $f : \mathbf{X} \rightarrow \mathbf{X}$ be an endomorphism in a rigid category. As above, it corresponds to a morphism $\check{f} : \mathbf{1} \rightarrow \check{\mathbf{X}} \otimes \mathbf{X}$. Composing with the commutativity followed by the evaluation we get a morphism $\text{Tr}_{\mathbf{X}}(f) : \mathbf{1} \rightarrow \mathbf{1}$ called the *trace* of f . The trace map $\text{Tr}_{\mathbf{X}} : \text{End}(\mathbf{X}) \rightarrow \mathbb{k} := \text{End}(\mathbf{1})$ is multiplicative (with respect to the tensor product), and $\text{Tr}_{\mathbf{1}}$ is the identity. The *rank* of an object $\mathbf{X}$ is defined to be $\text{Tr}_{\mathbf{X}}(\text{id}_{\mathbf{X}})$. Thus it is an element of $\mathbb{k}$.

3.3. A monoidal functor from a rigid category is automatically closed. If F is a monoidal functor from $\mathcal{C}$ to $\mathcal{D}$, where $\mathcal{C}$ is rigid, then $F(\mathbb{k}_{\mathcal{C}}) \subseteq \mathbb{k}_{\mathcal{D}}$, and in this sense, F preserves the trace. In particular, $\text{rk}(\mathbf{X}) = \text{rk}(F(\mathbf{X}))$.

3.4. Definition. A *tensor category* is a rigid monoidal category $\mathcal{C}$ which is abelian. It follows that $\otimes$ is additive and exact in each coordinate. It also follows that $\mathbb{k}$ is a commutative ring, and that $\mathcal{C}$ has a natural $\mathbb{k}$ -linear structure, where $\otimes$ is $\mathbb{k}$ -bilinear, and $\text{Tr}_{\mathbf{X}}$ is $\mathbb{k}$ -linear. A *tensor functor* is a monoidal functor between tensor categories which is additive (and therefore $\mathbb{k}$ -linear).

Given an object $\mathbf{X}$ of a tensor category $\mathcal{C}$, we denote by $\mathcal{C}_{\mathbf{X}}$ the full subcategory of $\mathcal{C}$ whose objects are isomorphic to sub-quotients of finite sums of tensor powers of $\mathbf{X}$ (including $\mathbf{X}^{\otimes -1} = \check{\mathbf{X}}$ and its powers). $\mathcal{C}_{\mathbf{X}}$ is a tensor sub-category of $\mathcal{C}$, and $\mathcal{C}$ is a filtered union of sub-categories of this form. $\mathbf{X}$ is called a *tensor generator* for $\mathcal{C}$ if $\mathcal{C} = \mathcal{C}_{\mathbf{X}}$. For most of what follows we will restrict our attention to categories that have a tensor generator.

3.5. Definition. A *neutral Tannakian category* over a field $\mathbb{k}$ is a tensor category $\mathcal{C}$ with $\text{End}(\mathbf{1}) = \mathbb{k}$, which admits an exact tensor functor into the category $\mathcal{V}ec_{\mathbb{k}}$ of finite-dimensional $\mathbb{k}$ -vector spaces (with the usual tensor structure). Such a functor is called a *fibre functor*. It is automatically faithful.

Given a fibre functor ω and a $\mathbb{k}$ -algebra A , the functor $\omega \otimes_{\mathbb{k}} A$ (with $(\omega \otimes_{\mathbb{k}} A)(\mathbf{X}) = \omega(\mathbf{X}) \otimes_{\mathbb{k}} A$) is again an exact tensor functor (into a tensor subcategory of the category $\mathcal{V}ec_A$ of projective finitely presented A -modules). We thus get a functor $A \mapsto \text{Aut}(\omega \otimes_{\mathbb{k}} A)$ from the category of $\mathbb{k}$ -algebras to groups. This functor is denoted by $\underline{\text{Aut}}_{\mathbb{k}}^{\otimes}(\omega)$.

3.6. Example. If $\mathcal{G}$ is an affine group scheme over $\mathbb{k}$, the category $\mathcal{R}ep_{\mathcal{G}}$ of (finite-dimensional) representations of $\mathcal{G}$ over $\mathbb{k}$ is an abelian tensor category which neutral Tannakian, as witnessed by the forgetful functor ω . If A is a $\mathbb{k}$ -algebra, and $g \in \mathcal{G}(A)$, the action of $\mathcal{G}$ exhibits g as an automorphism of $\omega \otimes_{\mathbb{k}} A$. This determines a map (of functors) $\mathcal{G} \rightarrow \underline{\text{Aut}}_{\mathbb{k}}^{\otimes}(\omega)$. The main theorem states that this map is an isomorphism, and that this example is the most general one:

3.7. Theorem (Saavedra). Let $\mathcal{C}$ be a neutral Tannakian category, and let $\omega : \mathcal{C} \rightarrow \mathcal{V}ec_{\mathbb{k}}$ be a fibre functor.

3.7.1. The functor $\underline{\text{Aut}}_{\mathbb{k}}^{\otimes}(\omega)$ is representable by an affine group scheme $\mathcal{G}$ over $\mathbb{k}$.

3.7.2. The fibre functor ω factors through a tensor equivalence $\omega : \mathcal{C} \rightarrow \mathcal{R}ep_{\mathcal{G}}$.

3.7.3. If $\mathcal{C} = \mathcal{R}ep_{\mathcal{H}}$ for some affine group scheme $\mathcal{H}$, then the natural map $\mathcal{H} \rightarrow \mathcal{G}$ given in 3.6 is an isomorphism.

In the following section we present a proof of (a slightly weaker version of) this theorem, using the model theoretic tools of section 2.

4. THE THEORY ASSOCIATED WITH A TENSOR CATEGORY

In this section, we associate a theory with any Tannakian category, and use theorem 2.5 to prove the main theorem 3.7.

4.1. Let $(\mathcal{C}, \otimes, \phi, \psi)$ be a tensor category over a field $\mathbb{k}$ of characteristic 0, such that the rank of every object is a natural number. Let K be an extension field of $\mathbb{k}$. The theories $T_{\mathcal{C}}$ and $\tilde{T}_{\mathcal{C}}$ (depending on K , which is omitted from the notation) are defined as follows:

4.1.1. $T_{\mathcal{C}}$ has a sort $\mathbf{L}$, as well as a sort $V_{\mathbf{X}}$ for every object $\mathbf{X}$ of $\mathcal{C}$, and a function symbol $v_f : V_{\mathbf{X}} \rightarrow V_{\mathbf{Y}}$ for any morphism $f : \mathbf{X} \rightarrow \mathbf{Y}$. There are a binary operation symbol $+_{\mathbf{X}}$ on each $V_{\mathbf{X}}$, and a function symbol $\cdot_{\mathbf{X}} : \mathbf{L} \times V_{\mathbf{X}} \rightarrow V_{\mathbf{X}}$ for each object $\mathbf{X}$. $\mathbf{L}$ contains a constant symbol for each element of K .

4.1.2. The theory says that $\mathbf{L}$ is an algebraically closed field, whose field operations are given by $+_1$ and $\cdot_1$. The restriction of the operation to the constant symbols is given by the field structure on K .

It also says that $+_{\mathbf{X}}$ and $\cdot_{\mathbf{X}}$ determine a vector space structure over $\mathbf{L}$ on every $V_{\mathbf{X}}$, and $V_{\mathbf{X}}$ has dimension $\text{rk}(\mathbf{X})$ over $\mathbf{L}$. Each v_f is an $\mathbf{L}$ -linear map.

4.1.3. The $\mathbb{k}$ -linear category structure is reflected in the theory: $v_{id_{\mathbf{X}}}$ is the identity map for each $\mathbf{X}$, $v_f = 0$ if and only if $f = 0$, $v_{f \circ g} = v_f \circ v_g$, if $a \in \mathbb{k}$ is viewed as an element of $\text{End}(\mathbf{X})$, then v_a is multiplication by $a \in K$. Also, $v_{f+g} = v_f + v_g$, and v_f is injective or surjective if and only if f is.

4.1.4. For any two objects $\mathbf{X}$ and $\mathbf{Y}$ there is a further function symbol $b_{\mathbf{X},\mathbf{Y}} : V_{\mathbf{X}} \times V_{\mathbf{Y}} \rightarrow V_{\mathbf{X} \otimes \mathbf{Y}}$. The theory says that $b_{\mathbf{X},\mathbf{Y}}$ is bilinear, and the induced map $V_{\mathbf{X}} \otimes_{\mathbf{L}} V_{\mathbf{Y}} \rightarrow V_{\mathbf{X} \otimes \mathbf{Y}}$ is an isomorphism. Here $\otimes_{\mathbf{L}}$ is the usual tensor product of vector spaces over $\mathbf{L}$ (in a given model). This statement is first order, since the spaces are finite-dimensional.

The theory also says that $v_{\psi_{\mathbf{X},\mathbf{Y}}}(b_{\mathbf{X},\mathbf{Y}}(x, y)) = b_{\mathbf{Y},\mathbf{X}}(y, x)$, and similarly for ϕ .

If $u \in V_{\mathbf{X}}$ and $v \in V_{\mathbf{Y}}$ (either terms or elements in some model), we write $u \otimes v$ for $b_{\mathbf{X},\mathbf{Y}}(u, v)$.

4.1.5. The theory $\widetilde{T}_{\mathcal{C}}$ is the expansion of $T_{\mathcal{C}}$ by an extra sort $P_{\mathbf{X}}$, for every object $\mathbf{X}$, together with a surjective map $\pi_{\mathbf{X}} : V_{\mathbf{X}} \rightarrow P_{\mathbf{X}}$ identifying $P_{\mathbf{X}}$ with the projective space associated with $V_{\mathbf{X}}$. It also includes function symbols $p_f : P_{\mathbf{X}} \rightarrow P_{\mathbf{Y}}$ and $d_{\mathbf{X},\mathbf{Y}} : P_{\mathbf{X}} \times P_{\mathbf{Y}} \rightarrow P_{\mathbf{X} \otimes \mathbf{Y}}$ for any objects $\mathbf{X}, \mathbf{Y}$ and morphism f of $\mathcal{C}$, and the theory says that they are the projectivisations of the corresponding maps v_f and $b_{\mathbf{X},\mathbf{Y}}$.

4.2. Proposition. The theory $T_{\mathcal{C}}$ is stable. In particular, $\mathbf{L}$ is stably embedded. In each model, $\mathbf{L}$ is a pure algebraically closed field (possibly with additional constants).

Proof. A choice of basis for each vector space identifies it with a power of $\mathbf{L}$. All linear and bilinear maps are definable in the pure field structure on $\mathbf{L}$. Since $\mathbf{L}$ is stable and stability is not affected by parameters, so is $T_{\mathcal{C}}$. $\square$

4.3. Proposition. The theories $T_{\mathcal{C}}$ and $\widetilde{T}_{\mathcal{C}}$ eliminate quantifiers (possibly after naming some constants). The theory $\widetilde{T}_{\mathcal{C}}$ eliminates imaginaries.

Proof. That $\widetilde{T}_{\mathcal{C}}$ eliminates imaginaries is precisely the statement of Hrushovski [12, prop. 4.2] (see also 5.5.6). The proof of quantifier elimination is similar: since all sorts are interpretable (with parameters) in ACF , all definable sets are boolean combinations of Zariski-closed subsets. A Zariski-closed subset of $V_{\mathbf{X}}$ or $P_{\mathbf{X}}$ is the set of zeroes of polynomials, which are elements of the symmetric algebra on $\widetilde{V}_{\mathbf{X}}$. But $\widetilde{V}_{\mathbf{X}}$ is identified with $V_{\widetilde{\mathbf{X}}}$, and the tensor algebra on it, as well as the action on $V_{\mathbf{X}}$ or $P_{\mathbf{X}}$ is definable without quantifiers. $\square$

4.4. Models and fibre functors. Models of $T_{\mathcal{C}}$ are essentially fibre functors over algebraically closed fields. More precisely, we have the following statements.

4.4.1. Let M be a model of $T_{\mathcal{C}}$, and let A be a subset of M such that for any object $\mathbf{X}$, $V_{\mathbf{X}}(A)$ contains a basis of $V_{\mathbf{X}}(M)$ over $\mathbf{L}(M)$. Then $\mathbf{X} \mapsto V_{\mathbf{X}}(A)$ determines a fibre functor over the field extension $\mathbf{L}(A)$ of K .

Proof. We first note that since (by definition) $\mathbf{L}(A)$ is definably closed, it is indeed a field. Likewise, since each $V_{\mathbf{X}}(A)$ is definably closed, it is a vector space over $\mathbf{L}(A)$, and the $V_{\mathbf{X}}(A)$ are closed under the linear maps that come from $\mathcal{C}$. Thus $\mathbf{X} \mapsto V_{\mathbf{X}}(A)$ defines a $\mathbb{k}$ -linear functor from $\mathcal{C}$ to the category of vector spaces over $\mathbf{L}(A)$.

The map $b_{\mathbf{X},\mathbf{Y}}$ determines a map $c_{\mathbf{X},\mathbf{Y}} : V_{\mathbf{X}}(A) \otimes_{\mathbf{L}(A)} V_{\mathbf{Y}}(A) \rightarrow V_{\mathbf{X} \otimes \mathbf{Y}}(A)$. Since the map $V_{\mathbf{X}}(A) \otimes_{\mathbf{L}(A)} V_{\mathbf{Y}}(A) \rightarrow V_{\mathbf{X}}(M) \otimes_{\mathbf{L}(M)} V_{\mathbf{Y}}(M)$ is injective, so is $c_{\mathbf{X},\mathbf{Y}}$, and so the whole statement will follow from knowing that the dimension of $V_{\mathbf{X}}(A)$ over $\mathbf{L}(A)$ is $\text{rk}(\mathbf{X})$. Since $V_{\mathbf{X}}(A)$ contains a basis over $\mathbf{L}(M)$, we know that it is at least $\text{rk}(\mathbf{X})$.

For any $a_1, \dots, a_n \in V_{\mathbf{X}}(A)$, the set of all $(x_1, \dots, x_n) \in \mathbf{L}^n$ such that $\sum x_i a_i = 0$ is an A -definable subspace of $\mathbf{L}^n$. Since $\mathbf{L}$ is stably embedded, it is $\mathbf{L}(A)$ definable. Hence it has an $\mathbf{L}(A)$ -definable basis (Milne [22, Lemma 4.10]). This shows that the dimension is correct. $\square$

4.4.2. Let A be a subset of a model M of $T_{\mathcal{C}}$ such that $\text{dcl}(A \cup \mathbf{L}(M)) = M$, and such that $\mathbf{L}(A)$ is algebraically closed. Then $\text{dcl}(A)$ is a model. In particular, $\mathbf{X} \mapsto V_{\mathbf{X}}(A)$ determines a fibre functor for $\mathcal{C}$ over $\mathbf{L}(A)$.

Proof. We only need to prove that each $V_{\mathbf{X}}(A)$ contains a basis over $\mathbf{L}(M)$. By assumption, there is an A -definable map f from a subset $\mathbf{U}$ of $\mathbf{L}^m$ to $V_{\mathbf{X}}^n$, where $n = \text{rk}(\mathbf{X})$, such that $f(u)$ is a basis of $V_{\mathbf{X}}(M)$ for all $u \in \mathbf{U}(M)$. Since $\mathbf{U}$ is $\mathbf{L}(A)$ -definable, and $\mathbf{L}(A)$ is algebraically closed, $\mathbf{U}(A)$ contains a point u , and $f(u)$ is thus a basis in $V_{\mathbf{X}}(A)$. $\square$

4.4.3. Conversely, if K_1 is a field extension of K , $\omega : \mathcal{C} \rightarrow \text{vec}_{K_1}$ is a fibre functor, and $\bar{K}$ is an algebraically closed field containing K_1 , the assignment $V_{\mathbf{X}} \mapsto \omega(\mathbf{X}) \otimes_{K_1} \bar{K}$ determines a model M_{ω} of $T_{\mathcal{C}}$, such that $A = \omega \subseteq M$ satisfies $\text{dcl}(A \cup \mathbf{L}(M_{\omega})) = M_{\omega}$, and $\mathbf{L}(A) = K_1$.

In particular, $T_{\mathcal{C}}$ is consistent if and only if $\mathcal{C}$ has a fibre functor.

Proof. Since $\bar{\omega} = \omega \otimes_{K_1} \bar{K}$ is itself a fibre functor, it is clear that M_{ω} is a model ($b_{\mathbf{X}, \mathbf{Y}}$ is defined to be the composition of the canonical pairing $\bar{\omega}(\mathbf{X}) \times \bar{\omega}(\mathbf{Y}) \rightarrow \bar{\omega}(\mathbf{X}) \otimes_{\bar{K}} \bar{\omega}(\mathbf{Y})$ with the structure map $c_{\mathbf{X}, \mathbf{Y}} : \bar{\omega}(\mathbf{X}) \otimes_{\bar{K}} \bar{\omega}(\mathbf{Y}) \rightarrow \bar{\omega}(\mathbf{X} \otimes \mathbf{Y})$). Also, it is clear that $\text{dcl}(A \cup \mathbf{L}(M_{\omega})) = M_{\omega}$. To prove that $\mathbf{L}(A) = K_1$, we note that any automorphism of $\bar{K}$ over K_1 extends to an automorphism of the model by acting the second term. $\square$

4.5. From now on we assume that $\mathcal{C}$ has a fibre functor ω , and denote the theory of the model M_{ω} constructed in 4.4.3 by T_{ω} (so T_{ω} is a completion of $T_{\mathcal{C}}$).

We also assume that the category $\mathcal{C}$ has a tensor generator (3.4). The general results are obtained as a limit of this case, in the usual way. (This is not really necessary, since, as mentioned in 2.4, the theory works in general, but it makes it easier to apply standard model theoretic results about definable, rather than pro-definable set.) We fix a tensor generator Q .

4.6. **Claim.** The theory $T_{\mathcal{C}}$ is an internal cover of $\mathbf{L}$, viewed as an interpretation of ACF_K .

Proof. $\mathbf{L}$ is stably embedded in $T_{\mathcal{C}}$ since $T_{\mathcal{C}}$ is stable. Given a model M of $T_{\mathcal{C}}$, any element of $V_{\mathbf{X}}(M)$ is a linear combination of elements of a basis a for $V_{\mathbf{X}}(M)$ over $\mathbf{L}(M)$, hence is definable over a and $\mathbf{L}(M)$. We note that if Q is a tensor generator, then a basis for V_Q determines a basis for any other $V_{\mathbf{X}}$ (This is clear for tensor products, direct sums, duals and quotients. For a sub-object $\mathbf{Y} \rightarrow \mathbf{X}$, the set of all tuples x in $\mathbf{L}$ such that $\sum x_i a_i \in V_{\mathbf{Y}}$ is a -definable, hence has a point in $\mathbf{L}(a)$.) $\square$

4.7. **Proof of 3.7.1.** Let $\mathbf{G}$ be the definable group in $T_{\mathcal{C}}$ corresponding to it as an internal cover of $\mathbf{L}$ (2.5). By 4.4.3, ω determines a definably closed subset A of an arbitrarily large model M , such that $\text{dcl}(A \cup \mathbf{L}(M)) = M$ and $\mathbf{L}(A) = \mathbb{k}$. Given a field extension K , we may assume that $\mathbf{L}(M)$ contains K .

For each object $\mathbf{X}$, $V_{\mathbf{X}}(A \cup K) = \omega(\mathbf{X}) \otimes_{\mathbb{k}} K$ (again by 4.4.3). It follows by 2.6 that for such field K , $\mathbf{G}(\text{dcl}(A \cup K)) = \text{Aut}(\text{dcl}(A \cup K)/K)$, and since by the

definition of the theory, such automorphisms are the same as automorphisms of $\omega \otimes_{\mathbb{k}} K$ as a tensor functor, we get that $\mathbf{G}(\mathrm{dcl}(A \cup K)) = \underline{\mathrm{Aut}}^{\otimes}(\omega)(K)$.

On the other hand, A satisfies assumption 2.7, hence we get a definable group $\mathbf{G}_A$ in $\mathbf{L}$ (i.e., in ACF), such that $\mathbf{G}_A(K) = \mathbf{G}(\mathrm{dcl}(A \cup K))$. By Poizat [26, p. 4.5], any such group is algebraic.

4.8. Remark. By inspecting the proof of 2.6 and the construction of $\mathbf{G}$ in this particular case, it is easy to extend the above proof to the case when K is an integral domain. However, I don't know how to use the same argument for more general K .

4.9. Proof of 3.7.3. Let $\mathcal{C} = \mathcal{R}ep_{\mathcal{H}}$, where $\mathcal{H}$ is an algebraic group over $\mathbb{k}$, which we view as a definable group $\mathbf{H}$ in $ACF_{\mathbb{k}}$. Let $\mathbf{G}$ be the definable internality group in T_{ω} and A the definable subset corresponding to the (forgetful) fibre functor ω , all as in 4.7. We obtain a definable group $\mathbf{G}_A$, and the action of $\mathcal{H}$ on its representations determines a map of group functors from $\mathbf{H}$ to $\mathbf{G}_A$, which by Beth definability is definable and therefore algebraic. This homomorphism is injective (on points in any field extension) since $\mathbf{H}$ has a faithful representation. We identify $\mathbf{H}$ with its image in $\mathbf{G}_A$.

By the Galois correspondence (2.5), to prove that $\mathbf{H} = \mathbf{G}_A$ it is enough to show that any A -definable element in $T_{\mathcal{C}}^{eq}$ fixed by $\mathbf{H}(M)$ is also fixed by $\mathbf{G}_A(M)$, i.e., is 0-definable. By 4.3, $T_{\mathcal{C}}^{eq} = \tilde{T}_{\mathcal{C}}$. If $v \in V_{\mathbf{X}}(A)$ is fixed by $\mathbf{H}$, then the map $\mathbf{L} \rightarrow V_{\mathbf{X}}$ given by $1 \mapsto v$ is a map of $\mathbf{H}$ representations, hence is given by a function symbol f , so $v = f(1)$ is 0-definable. Likewise, if $p \in P_{\mathbf{X}}(A)$ is fixed by $\mathbf{H}$, then it corresponds to a sub-representation $l_p \subset V_{\mathbf{X}}$ (over $\mathbb{k}$), hence is given by a predicate, so again p is 0-definable.

This proves that the map from $\mathbf{H}$ to $\mathbf{G}_A$ is bijective on points in any field. Any such algebraic map (in characteristic 0) is an isomorphism (Waterhouse [30, p. 11.4]).

4.10. Proof of 3.7.2 (sketch). Let $T_{\mathcal{C}}$, $\mathbf{G}$, A and $\mathbf{G}_A$ be as above, and let $\mathcal{V}ec_{\mathbf{L}}$ be the category of definable $\mathbf{L}$ -vector spaces interpretable in $T_{\mathcal{C}}$, with definable linear maps between them as morphisms. This is clearly a tensor category, and $\mathbf{X} \mapsto V_{\mathbf{X}}$ is an exact faithful $\mathbb{k}$ -linear tensor functor from $\mathcal{C}$ to $\mathcal{V}ec_{\mathbf{L}}$.

On the other hand, a representation of $\mathbf{G}_A$ is given by some definable action in ACF , and by 2.8, this action comes from some action of $\mathbf{G}$ on a definable set in $T_{\mathcal{C}}$, which must therefore be a vector space over $\mathbf{L}$. Conversely, each linear action of $\mathbf{G}$ on an object in $\mathcal{V}ec_{\mathbf{L}}$ gives a representation of $\mathbf{G}_A$ after taking A -points, and similarly for morphisms. Thus, we get an equivalence (of $\mathbb{k}$ -linear tensor categories) between $\mathcal{V}ec_{\mathbf{L}}$ and $\mathcal{R}ep_{\mathbf{G}_A}$.

It remains to show that the functor $\mathbf{X} \mapsto V_{\mathbf{X}}$ from $\mathcal{C}$ to $\mathcal{V}ec_{\mathbf{L}}$ is full and surjective on isomorphism classes, i.e., that any definable $\mathbf{L}$ -vector space is essentially of the form $V_{\mathbf{X}}$, and that any map between is of the form v_f . To prove the first, we consider the group $\mathbf{G}$ and its torsor $\mathbf{X}$. These are defined by some 0-definable subspaces of powers of V_Q (where Q is a tensor generator.) By examining the explicit definition of $\mathbf{G}$ and $\mathbf{X}$, one concludes that these 0-definable subspaces in fact come from $\mathcal{C}$. We omit the proof, since it is very similar to the original algebraic one. Once this is known, given any other representation $\mathbf{D}$, the set $\mathbf{X}_{\mathbf{D}}$ in 2.8 is explicitly given by tensor operations, and so comes from $\mathcal{C}$ as well. Then proof for morphisms is again similar.

4.11. Corollary. Quantifier elimination in $T_{\mathcal{C}}$ holds without parameters: any definable set is quantifier-free without parameters.

Proof. This holds anyway with parameters from the definable closure of the empty set. We now know that any such parameter is given by a term in the language. $\square$

4.12. Remark. In Hrushovski [12], an equivalence is constructed between internal covers of a complete theory $\mathcal{T}$, and connected definable groupoids in $\mathcal{T}$. It follows from theorem 2.8 there that if $T_{\mathcal{C}}$ is consistent and the induced structure on $\mathbf{L}$ is precisely $ACF_{\mathbb{k}}$, then the associated groupoid is connected. It also follows that $T_{\mathcal{C}}$ itself is complete. It then follows from lemma 2.4 in the same paper that any two models of $T_{\mathcal{C}}$ are isomorphic. Given two fibre functors of $\mathcal{C}$ (over some extension field), extending both to a model of $T_{\mathcal{C}}$ thus shows that the fibre functors are locally isomorphic. Conversely, if $T_{\mathcal{C}}$ induces new structure on $\mathbf{L}$, it is easy to construct fibre functors which are not locally isomorphic (cf. Deligne and Milne [5, p. 3.15]).

The main result of Deligne [4] can be viewed as stating that the induced structure on $\mathbf{L}$ is indeed $ACF_{\mathbb{k}}$. More precisely, given a fibre functor ω over an (affine) scheme $\mathcal{S}$ over $\mathbb{k}$, Deligne constructs a groupoid scheme $\underline{Aut}_{\mathbb{k}}^{\otimes}(\omega)$, with object scheme $\mathcal{S}$. The main result of Deligne [4] (theorem 1.12) states that this groupoid scheme is faithfully flat over $\mathcal{S} \times \mathcal{S}$ (hence connected). A K -valued point of $\mathcal{S}$ (for some field K) determines a fibre functor ω_K over K . Viewing this fibre functor as subset of a model, a choice of a basis for a tensor generator determines a type over $\mathbf{L}$, and therefore an object of the groupoid corresponding to $T_{\mathcal{C}}$ (choosing a different basis amounts to choosing a different object which is isomorphic over K in the groupoid). It is clear from definition of $\underline{Aut}_{\mathbb{k}}^{\otimes}(\omega)$ that this process gives an equivalence from this groupoid to the groupoid associated with the internal cover $T_{\mathcal{C}}$. Thus, Deligne's theorem implies that this groupoid is connected, and therefore that the induced structure on $\mathbf{L}$ is that of $ACF_{\mathbb{k}}$. It seems plausible that there should be a direct model theoretic proof of this result, but I could not find it. We note that the non-existence of new structure also directly implies (by the existence of prime models) Deligne [4, corollary 6.20], which states that $\mathcal{C}$ has a fibre functor over the algebraic closure of $\mathbb{k}$.

We note also that any definable groupoid in $ACF_{\mathbb{k}}$ is equivalent to a groupoid scheme (any such groupoid is equivalent to one with a finite set of objects, and the automorphism group of each object is algebraic). The category $\mathcal{C}$ of representations of the (gerb associated with the) groupoid is a tensor category satisfying $End(\mathbf{1}) = \mathbb{k}$ (Deligne and Milne [5, section 3]), and it is easy to see (using the same methods as in 4.9) that the groupoid associated with the internal cover $T_{\mathcal{C}}$ is equivalent to the original one. Therefore, we obtain that any internal cover of $ACF_{\mathbb{k}}$ has (up to equivalence) the form $T_{\mathcal{C}}$ for some Tannakian category over $\mathbb{k}$.

4.13. Remark. We interpret model theoretically two additional result of Deligne [4]. In section 7, it is shown that in our context (i.e., $char(\mathbb{k}) = 0$ and $rk(\mathbf{X})$ is a natural number), $\mathcal{C}$ always has a fibre functor. In light of 4.4.3, this result asserts that $T_{\mathcal{C}}$ is consistent for any such $\mathcal{C}$.

In section 8, Deligne defines, for a Tannakian category $\mathcal{C}$ (and even somewhat more generally), a $\mathcal{C}$ -group $\pi(\mathcal{C})$, called the *fundamental group* of $\mathcal{C}$ (a $\mathcal{C}$ -group is a commutative Hopf algebra object in $\text{Ind}(\mathcal{C})$; similarly for affine $\mathcal{C}$ -schemes). The basic idea is that the concepts associated with neutral Tannakian categories, also

make sense for “fibre functors” into other tensor categories, and $\pi(\mathcal{C})$ is obtained by reconstructing a group from the identity functor.

Thus, $\pi(\mathcal{C})$ comes with an action on each object, commuting with all morphisms. It thus represents the tensor automorphisms of the identity functor from $\mathcal{C}$ to itself, in the sense that for any affine $\mathcal{C}$ -scheme $\mathbf{X}$, the action identifies $\pi(\mathcal{C})(\mathbf{X})$ with the group of tensor automorphisms of the functor $\mathbf{A} \mapsto \mathbf{A} \otimes \mathbf{X}$ (from $\mathcal{C}$ to the tensor category of vector bundles over $\mathbf{X}$). It has the property that for any “real” fibre functor ω over a scheme $\mathcal{S}$, applying ω to the action identifies $\omega(\pi(\mathcal{C}))$ with the group scheme $\underline{\text{Aut}}_{\mathcal{S}}^{\otimes}(\omega)$ constructed.

Using either the explicit construction, or the properties above, it is clear that in terms of the theory $T_{\mathcal{C}}$, $\pi(\mathcal{C})$ is nothing but the definable automorphism group (more precisely, the Hopf algebra object that defines it maps to an ind-definable Hopf algebra in $T_{\mathcal{C}}$, which is definably isomorphic to the ind-definable Hopf algebra of functions on the internality group). This also compares with 1.13: the group G is obtained by considering the automorphisms of (essentially) the identity functor, rather than of the “fibre functor” F there.

4.14. Remark. One can recover in a similar manner (and somewhat more easily) Grothendieck’s approach to usual Galois theory (cf. Grothendieck et al. [8, Exposé V.4]). Briefly, given a category $\mathcal{C}$ with a “fibre functor” F into the category of finite sets, satisfying conditions (G1)–(G6), one constructs as above a theory $T_{\mathcal{C}}$ with a sort $V_{\mathbf{X}}$ for each object, and a function symbol for each morphism. $T_{\mathcal{C}}$ is then the theory of F viewed as a structure in this language. Since every sort is finite, they are all internal to $\mathbf{2}$ (the co-product of the terminal object $\mathbf{1}$ with itself). Conditions (G1)–(G5) ensure that any definable set in fact comes from $\mathcal{C}$, and (G2), (G5) ensure that $T_{\mathcal{C}}$ has elimination of imaginaries. The Galois objects P_i that appear in the proof are precisely the 1-types of internality parameters that appear in the model theoretic construction of Galois group.

5. DIFFERENTIAL TANNAKIAN CATEGORIES

Our purpose in this section is to define differential tensor categories, and to give a model theoretic proof of the basic theorem, corresponding such categories, endowed with a suitably defined fibre functor, with linear differential algebraic groups. The method is completely analogous to that in the previous section.

Throughout this section, $\mathbb{k}$ is a field of characteristic 0.

5.1. Prolongations of abelian categories. We assume that in a tensor category $(\mathcal{C}, \otimes)$, the functor $\otimes$ is exact; this is automatic if $\mathcal{C}$ is rigid (see Deligne and Milne [5, prop. 1.16].)

5.1.1. Definition. Let $\mathcal{C}$ be a $\mathbb{k}$ -linear category. The *prolongation* $\mathcal{P}(\mathcal{C})$ of $\mathcal{C}$ is defined as follows: The objects are exact sequences $\mathbf{X} := 0 \rightarrow \mathbf{X}_0 \xrightarrow{i_{\mathbf{X}}} \mathbf{X}_1 \xrightarrow{\pi_{\mathbf{X}}} \mathbf{X}_0 \rightarrow 0$ of $\mathcal{C}$, and the morphisms between such objects are morphisms of exact sequences whose two $\mathbf{X}_0$ parts coincide.

An exact functor $F : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ gives rise to an induced functor $\mathcal{P}(F) : \mathcal{P}(\mathcal{C}_1) \rightarrow \mathcal{P}(\mathcal{C}_2)$. We denote by Π_i ($i = 0, 1$) the functors from $\mathcal{P}(\mathcal{C})$ to $\mathcal{C}$ assigning $\mathbf{X}_i$ to the object $0 \rightarrow \mathbf{X}_0 \xrightarrow{i_{\mathbf{X}}} \mathbf{X}_1 \xrightarrow{\pi_{\mathbf{X}}} \mathbf{X}_0 \rightarrow 0$ of $\mathcal{P}(\mathcal{C})$ (thus there is an exact sequence $0 \rightarrow \Pi_0 \xrightarrow{i_{\Pi}} \Pi_1 \xrightarrow{\pi_{\Pi}} \Pi_0 \rightarrow 0$). $\Pi_i(\mathbf{X})$ is also abbreviated as $\mathbf{X}_i$, and $\mathbf{X}$ is said to be over $\mathbf{X}_0$ (and similarly for morphisms.)

We note that $\mathcal{P}(\mathcal{C})$ can be viewed as the full subcategory of the category of “differential objects” in $\mathcal{C}$, consisting of objects whose homology is 0. A differential object is a pair $(\mathbf{X}, \phi)$ where $\mathbf{X}$ is an object of $\mathcal{C}$ and ϕ is an endomorphism of $\mathbf{X}$ with $\phi^2 = 0$. A morphism is a morphism in $\mathcal{C}$ that commutes with ϕ , and the homology is $\ker(\phi)/\text{im}(\phi)$. This is the same as the category of $\mathbb{k}[\epsilon]$ -modules in $\mathcal{C}$, in the sense of Deligne and Milne [5, p. 155] (where $\epsilon^2 = 0$). The advantage of this category is that it is again $\mathbb{k}$ -linear. However, I don’t know how to extend the tensor structure (defined below) to this whole category (in particular, the tensor structure defined there does not seem to coincide with ours).

5.1.2. Let $\mathbf{A}$ and $\mathbf{B}$ be two objects over $\mathbf{X}_0$. Their *Yoneda sum* $\mathbf{A} \star \mathbf{B}$ is a new object over $\mathbf{X}_0$, defined as follows (this is the addition in Yoneda’s description of $\text{Ext}^1(\mathbf{X}_0, \mathbf{X}_0)$): the combined map $\mathbf{X}_0 \times \mathbf{X}_0 \rightarrow \mathbf{A}_1 \times \mathbf{B}_1$ factors through $\mathbf{A}_1 \times_{\mathbf{X}_0} \mathbf{B}_1$, and together with the map $\mathbf{X}_0 \xrightarrow{1, -1} \mathbf{X}_0 \times \mathbf{X}_0$ gives rise to a map $f : \mathbf{X}_0 \rightarrow \mathbf{A}_1 \times_{\mathbf{X}_0} \mathbf{B}_1$. Let $\mathbf{W}_1$ be the co-kernel of this map. The map f composed with the projection from $\mathbf{A}_1 \times_{\mathbf{X}_0} \mathbf{B}_1$ to $\mathbf{X}_0$ is 0, so we obtain an induced map $p : \mathbf{W}_1 \rightarrow \mathbf{X}_1$. The diagonal inclusion Δ of $\mathbf{X}_0$ in $\mathbf{W}_1$ together with p give rise to an exact sequence $0 \rightarrow \mathbf{X}_0 \xrightarrow{\Delta} \mathbf{W}_1 \xrightarrow{p} \mathbf{X}_0 \rightarrow 0$, which is the required object.

For any object $\mathbf{A}$ of $\mathcal{P}(\mathcal{C})$, we denote by $T(\mathbf{A})$ the object obtained by negating all arrows that appear in $\mathbf{A}$.

5.1.3. Let $(\mathcal{C}, \otimes, \phi_0, \psi_0)$ be a tensor category. An object $\mathbf{X}_0$ of $\mathcal{C}$ gives rise to a functor from $\mathcal{P}(\mathcal{C})$ to itself, by tensoring the exact sequence pointwise. Since we assumed $\otimes$ to be exact, this functor also commutes with Yoneda sums: $(\mathbf{A} \star \mathbf{B}) \otimes \mathbf{X}_0$ is canonically isomorphic with $(\mathbf{A} \otimes \mathbf{X}_0) \star (\mathbf{B} \otimes \mathbf{X}_0)$. Also, $T(\mathbf{A}) \otimes \mathbf{X}_0$ is isomorphic to $T(\mathbf{A} \otimes \mathbf{X}_0)$.

We endow $\mathcal{P}(\mathcal{C})$ with a monoidal structure. The tensor product $\mathbf{A} \otimes \mathbf{B}$ of the two $\mathcal{P}(\mathcal{C})$ objects $\mathbf{A}$ and $\mathbf{B}$ is defined as follows: After tensoring the first with $\mathbf{B}_0$ and the second with $\mathbf{A}_0$, we obtain two objects over $\mathbf{A}_0 \otimes \mathbf{B}_0$. We now take their Yoneda sum.

We shall make use of the following exact sequence.

5.1.4. *Lemma.* for any two objects $\mathbf{A}$ and $\mathbf{B}$ of $\mathcal{P}(\mathcal{C})$, there is an exact sequence

$$0 \rightarrow (\mathbf{A} \otimes T(\mathbf{B}))_1 \xrightarrow{i} \mathbf{A}_1 \otimes \mathbf{B}_1 \xrightarrow{\pi} (\mathbf{A} \otimes \mathbf{B})_1 \rightarrow 0 \quad (3)$$

where π is the quotient of the map obtained from the maps $\pi_{\mathbf{A}} \otimes 1$ and $1 \otimes \pi_{\mathbf{B}}$, and i is the restriction of the map obtained from the maps $i_{\mathbf{A}} \otimes 1$ and $-1 \otimes i_{\mathbf{B}}$.

Proof. Exactness in the middle follows directly from the definitions. We prove that π is surjective, the injectivity of i being similar. We shall use the Mitchell embedding theorem (cf. Freyd [6]), which reduces the question to the case of abelian groups.

We in fact prove that already the map

$$\mathbf{A}_1 \otimes \mathbf{B}_1 \xrightarrow{\pi} \mathbf{A}_0 \otimes \mathbf{B}_1 \times_{\mathbf{A}_0 \otimes \mathbf{B}_0} \mathbf{A}_1 \otimes \mathbf{B}_0 =: U$$

is surjective. Let y be an element of U , and let y_1 and y_2 be its two projections to the components of U . Since the map $\mathbf{A}_1 \otimes \mathbf{B}_1 \xrightarrow{\pi_{\mathbf{A}} \otimes 1} \mathbf{A}_0 \otimes \mathbf{B}_1$ is surjective, y_1 can be lifted to an element $\tilde{y}_1$ of $\mathbf{A}_1 \otimes \mathbf{B}_1$. We have that

$$(\pi_{\mathbf{A}} \otimes 1)((1 \otimes \pi_{\mathbf{B}})(\tilde{y}_1)) = (1 \otimes \pi_{\mathbf{B}})((\pi_{\mathbf{A}} \otimes 1)(\tilde{y}_1)) = (1 \otimes \pi_{\mathbf{B}})(y_1) = (\pi_{\mathbf{A}} \otimes 1)(y_2)$$

Let $z = (1 \otimes \pi_{\mathbf{B}})(\tilde{y}_1) - y_2$. Since z is killed by $\pi_{\mathbf{A}} \otimes 1$, it comes from an element, also z , of $\mathbf{A}_0 \otimes \mathbf{B}_0$. Let $\tilde{z}$ be a lifting of z to $\mathbf{A}_0 \otimes \mathbf{B}_1$, and denote by $\tilde{z}$ also its image in $\mathbf{A}_1 \otimes \mathbf{B}_1$ under the inclusion $i_{\mathbf{A}} \otimes 1$. Then $\tilde{y}_1 - \tilde{z}$ is a lifting of y . $\square$

5.1.5. Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be three objects of $\mathcal{P}(\mathcal{C})$. The associativity constraint ϕ_0 of $\mathcal{C}$ gives rise to an isomorphism of $(\mathbf{A} \otimes \mathbf{B}) \otimes \mathbf{C}$ with the quotient of

$$\mathbf{A}_1 \otimes \mathbf{B}_0 \otimes \mathbf{C}_0 \times_{\mathbf{A}_0 \otimes \mathbf{B}_0 \otimes \mathbf{C}_0} \mathbf{A}_0 \otimes \mathbf{B}_1 \otimes \mathbf{C}_0 \times_{\mathbf{A}_0 \otimes \mathbf{B}_0 \otimes \mathbf{C}_0} \mathbf{A}_0 \otimes \mathbf{B}_0 \otimes \mathbf{C}_1$$

that identifies the three natural inclusions of $\mathbf{A}_0 \otimes \mathbf{B}_0 \otimes \mathbf{C}_0$, and similarly for $\mathbf{A} \otimes (\mathbf{B} \otimes \mathbf{C})$. We thus get an associativity constraint ϕ on $\mathcal{P}(\mathcal{C})$, over ϕ_0 .

Likewise, the commutativity constraint ψ_0 induces a commutativity constraint ψ on $\mathcal{P}(\mathcal{C})$ over ψ_0 .

5.1.6. *Proposition.* The data $(\mathcal{P}(\mathcal{C}), \otimes, \phi, \psi)$ as defined above forms a symmetric monoidal category, and Π_0 is a monoidal functor. It is rigid if $\mathcal{C}$ is rigid.

Proof. We define the additional data. Verification of the axioms reduces, as in lemma 5.1.4, to the case of abelian groups, where it is easy.

Let $u : \mathbf{1}_0 \rightarrow \mathbf{1}_0 \otimes \mathbf{1}_0$ be an identity object of $\mathcal{C}$. We set $\mathbf{1} = 0 \rightarrow \mathbf{1}_0 \rightarrow \mathbf{1}_0 \oplus \mathbf{1}_0 \rightarrow \mathbf{1}_0 \rightarrow 0$. For any object $\mathbf{A}$ of $\mathcal{P}(\mathcal{C})$, $\mathbf{1} \otimes \mathbf{A}$ is identified via u with $0 \rightarrow \mathbf{A}_0 \rightarrow (\mathbf{A}_1 \times_{\mathbf{A}_0} (\mathbf{A}_0 \oplus \mathbf{A}_0)) / \mathbf{A}_0 \rightarrow \mathbf{A}_0 \rightarrow 0$. This is canonically isomorphic (over $\mathcal{C}$) to $\mathbf{A}$, and so $\mathbf{1}$ acquires a structure of an identity object.

Assume that $\mathcal{C}$ is rigid. For an object $\mathbf{A}$ of $\mathcal{P}(\mathcal{C})$, we set $\check{\mathbf{A}}$ to be the dual exact sequence $0 \rightarrow \check{\mathbf{A}}_0 \xrightarrow{i_{\check{\mathbf{A}}}} \check{\mathbf{A}}_1 \xrightarrow{\pi_{\check{\mathbf{A}}}} \check{\mathbf{A}}_0 \rightarrow 0$. We define an evaluation map $\mathbf{A} \otimes \check{\mathbf{A}} \rightarrow \mathbf{1}$ as follows: We need to define two maps from $\mathbf{A}_0 \otimes \check{\mathbf{A}}_1 \times \mathbf{A}_1 \otimes \check{\mathbf{A}}_0$ to $\mathbf{1}_0$, that agree on the two inclusions of $\mathbf{A}_0 \otimes \check{\mathbf{A}}_0$, and such that the resulting map restricts to the evaluation on $\mathbf{A}_0 \otimes \check{\mathbf{A}}_0$.

To construct the first map, we consider the exact sequence (3), for $\mathbf{B} = \check{\mathbf{A}}$. We claim that the evaluation map on $\mathbf{A}_1 \otimes \check{\mathbf{A}}_1$ restricts to 0 when composed with i . To prove this, it is enough to show that the pair of maps obtained from $ev_{\mathbf{A}_1}$ by composition with $i_{\mathbf{A}} \otimes 1$ and $-1 \otimes i_{\check{\mathbf{A}}}$ comes from a map $\mathbf{A}_0 \otimes \check{\mathbf{A}}_0 \rightarrow \mathbf{1}_0$. However, under the adjunction, this pair of maps corresponds to $(i_{\mathbf{A}}, -\pi_{\mathbf{A}})$, and so comes from the identity map on $\mathbf{A}_0$. It follows that $ev_{\mathbf{A}_1}$ induces a map on $(\mathbf{A} \otimes \check{\mathbf{A}})_1$, which is the required map. The second map is obtained by projecting to $\mathbf{A}_0 \otimes \check{\mathbf{A}}_0$, and using the evaluation map on $\mathbf{A}_0$. By definition, this second map commutes with the projections to $\mathbf{A}_0 \otimes \check{\mathbf{A}}_0$ and the second coordinate of $\mathbf{1}$, restricting to the evaluation on $\mathbf{A}_0$. To prove that the first map restricts to the evaluation as well, we note that there is a commutative diagram

$$\begin{array}{ccc} (\mathbf{A} \otimes \check{\mathbf{A}})_1 & \xrightarrow{i} & \mathbf{A}_1 \otimes \check{\mathbf{A}}_1 \\ \downarrow \pi_{\mathbf{A} \otimes \check{\mathbf{A}}} & & \downarrow \pi \\ \mathbf{A}_0 \otimes \check{\mathbf{A}}_0 & \xrightarrow{i_{\mathbf{A} \otimes \check{\mathbf{A}}}} & (\mathbf{A} \otimes \check{\mathbf{A}})_1 \end{array}$$

where i is the (restriction of the) map obtained from the two maps $i_{\mathbf{A}} \otimes 1$ and $1 \otimes i_{\check{\mathbf{A}}}$.

Since $\pi_{\mathbf{A} \otimes \tilde{\mathbf{A}}}$ is surjective, it is therefore enough to prove that the maps $ev_{\mathbf{A}_1} \circ i$ and $ev_{\mathbf{A}_0} \circ \pi_{\mathbf{A}}$ coincide. This is indeed the case, since they both correspond to the inclusion of $\mathbf{A}_0$ in $\mathbf{A}_1$. $\square$

5.2. Differential tensor categories.

5.2.1. *Definition.* A *differential structure* on a tensor category $\mathcal{C}$ is a tensor functor D from $\mathcal{C}$ to $\mathcal{P}(\mathcal{C})$ which is a section of Π_0 . If D_1 and D_2 are two differential structures on $\mathcal{C}$, a morphism from D_1 to D_2 is a morphism of tensor functors that induces the identity morphism under Π_0 . A *differential tensor category* is a tensor category together with a differential structure.

Let D be a differential structure on $\mathcal{C}$. Since D is a section of Π_0 , it is determined by $\partial = \Pi_1 \circ D$. In other words, on the abelian level, it is given by a functor $\partial : \mathcal{C} \rightarrow \mathcal{C}$, together with an exact sequence $0 \rightarrow Id \rightarrow \partial \rightarrow Id \rightarrow 0$. However, this description does not include the tensor structure. We also note that ∂ is necessarily exact.

5.2.2. Let $(\mathcal{C}, D)$ be a differential tensor category, let $\partial = \Pi_1 \circ D$, and let $A = End(\mathbf{1})$. Recall that for any object $\mathbf{X}$, $End(\mathbf{X})$ is an A -algebra. The functor ∂ defines another ring homomorphism $\partial_1 : A \rightarrow End(\partial(\mathbf{1}))$. Given $a \in A$, the morphism $\partial_1(a) - a$ in $End(\partial(\mathbf{1}))$ restricts to 0 on $\mathbf{1}$, and thus induces a morphism from $\mathbf{1}$ to $\partial(\mathbf{1})$. Similarly, its composition with the projection $\partial(\mathbf{1}) \rightarrow \mathbf{1}$ is 0, so it factors through $\mathbf{1}$. We thus get a new element a' of A .

Claim. *The map $a \mapsto a'$ is a derivation on A .*

Proof. We need to show that given elements $a, b \in A$, the maps $\partial(ab) - ab$ and $(\partial(a) - a)b + a(\partial(b) - b)$ coincide on $\mathbf{1}$. This follows from the formula $\partial(ab) - ab = \partial(a)(\partial(b) - b) + (\partial(a) - a)b$, together with the fact that $\partial(a)(\partial(b) - b)$ induces $a(\partial(b) - b)$ on $\mathbf{1}$. $\square$

5.2.3. *Example.* Let $\mathcal{C}$ be the tensor category $\mathcal{V}_{ec\mathbb{k}}$ of finite dimensional vector spaces over a field $\mathbb{k}$. Given a derivative $'$ on $\mathbb{k}$, we construct a differential structure on $\mathcal{C}$ as follows: For a vector space $\mathbf{X}$, define $d(\mathbf{X}) = \mathfrak{D} \hat{\otimes} \mathbf{X}$, where $\mathfrak{D}$ is the vector space with basis $1, \partial$, and $\hat{\otimes}$ is the tensor product with respect to the *right* vector space structure on $\mathfrak{D}$, given by $1 \cdot a = a \cdot 1$ and $\partial \cdot a = a' \cdot 1 + a \cdot \partial$. The exact sequence $D(\mathbf{X})$ is defined by $x \mapsto 1 \hat{\otimes} x$, $1 \hat{\otimes} x \mapsto 0$ and $\partial \hat{\otimes} x \mapsto x$, for any $x \in \mathbf{X}$. If $T : \mathbf{X} \rightarrow \mathbf{Y}$ is a linear map, $d(T) = 1 \hat{\otimes} T$. We shall write x for $1 \hat{\otimes} x$ and ∂x for $\partial \hat{\otimes} x$. The structure of a tensor functor is obtained by sending $\partial(x \otimes y) \in d(\mathbf{X} \otimes \mathbf{Y})$ to the image of $\partial(x) \otimes y \oplus x \otimes \partial(y)$ in $(D(\mathbf{X}) \otimes D(\mathbf{Y}))_1$.

5.2.4. *Claim.* The constructions in 5.2.3 and in 5.2.2 give a bijective correspondence between derivatives on $\mathbb{k}$ and isomorphism classes of differential structures on $\mathcal{V}_{ec\mathbb{k}}$.

Proof. If D_1 and D_2 are two differential structures, then $D_1(\mathbf{1}_0)$ and $D_2(\mathbf{1}_0)$ are both identity objects, and are therefore canonically isomorphic to the same object $\mathbf{1}$. If D_1 and D_2 are isomorphic, then the maps $d_i : End(\mathbf{1}_0) \rightarrow End(\mathbf{1})$ are conjugate, and therefore equal, since $End(\mathbf{1})$ is commutative.

It is clear from the definition that the derivative on $\mathbb{k}$ obtained from the differential structure associated with a derivative is the original one. Conversely, if D_1 and D_2 are two differential structures that give the same derivative on $\mathbb{k}$, then we may identify $D_1(\mathbf{1}_0)$ and $D_2(\mathbf{1}_0)$. Under this identification, we get that the maps d_i are the same. But the functors D_i are determined by d_i . $\square$

5.2.5. We now come to the definition of functors between differential tensor categories. For simplicity, we shall only define (and use) *exact* such functors.

Let $\omega : \mathcal{C} \rightarrow \mathcal{D}$ be an exact functor between abelian categories. There is an induced functor $\mathcal{P}(\omega) : \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{D})$, given by applying ω to each term. If $\mathcal{C}$ and $\mathcal{D}$ are tensor categories, the structure of a tensor functor on ω gives rise to a similar structure on $\mathcal{P}(\omega)$ (again, since ω is exact.) If $t : \omega_1 \rightarrow \omega_2$ is a (tensor) morphism of functors, we likewise get an induced morphism $\mathcal{P}(t) : \mathcal{P}(\omega_1) \rightarrow \mathcal{P}(\omega_2)$.

5.2.6. *Definition.* Let $(\mathcal{C}_1, D_1)$ and $(\mathcal{C}_2, D_2)$ be two differential tensor categories. A *differential tensor functor* from $\mathcal{C}_1$ to $\mathcal{C}_2$ is an exact tensor functor ω from $\mathcal{C}_1$ to $\mathcal{C}_2$, together with an isomorphism of tensor functors $r : \mathcal{P}(\omega) \circ D_1 \rightarrow D_2 \circ \omega$, such that the “restriction” $\Pi_0 \odot r : \omega \rightarrow \omega$, obtained by composing with Π_0 on $\mathcal{P}(\mathcal{C}_2)$, is the identity.

A morphism between two such differential tensor functors (ω_1, r_1) and (ω_2, r_2) is a morphism t between them as tensor functors such that the following diagram (of tensor functors and tensor maps between them) commutes:

$$\begin{array}{ccc} \mathcal{P}(\omega_1) \circ D_1 & \xrightarrow{r_1} & D_2 \circ \omega_1 \\ \downarrow \mathcal{P}(t) \odot D_1 & & \downarrow D_2 \odot t \\ \mathcal{P}(\omega_2) \circ D_1 & \xrightarrow{r_2} & D_2 \circ \omega_2 \end{array} \quad (4)$$

where $D_2 \odot t$ is the map from $D_2 \circ \omega_1$ to $D_2 \circ \omega_2$ obtained by applying D_2 to t “pointwise”.

Given a differential tensor functor ω , we denote by $\underline{\text{Aut}}^\partial(\omega)$ the group of automorphisms of ω .

If $\mathcal{C}$ is a differential tensor category, and $\mathbb{k} = \text{End}(\mathbf{1})$ is a field, a $\mathbb{k}$ -linear differential tensor functor into $\mathcal{V}_{ec\mathbb{k}}$ (with the induced differential structure) is called a *differential fibre functor*. Given such a functor ω , we denote by G_ω the functor from differential $\mathbb{k}$ -algebras to groups assigning to an algebra A the group $\underline{\text{Aut}}^\partial(A \otimes \omega)$.

5.2.7. Given a $\mathbb{k}$ vector space V , the map $d : V \rightarrow \mathfrak{D} \hat{\otimes} V$ given by $v \mapsto \partial v$ is a derivation, in the sense that $d(av) = a'v + ad(v)$ (where V is identified with its image in $\mathfrak{D} \hat{\otimes} V$.) It is universal for this property: any pair $(i, d) : V \rightarrow W$, where i is linear, and d is a derivation with respect to i factors through it.

Therefore, a fibre functor on $(\mathcal{C}, D)$ is a fibre functor ω in the sense of tensor categories, together with a functorial derivation $d_{\mathbf{X}_0} : \omega(\mathbf{X}_0) \rightarrow \omega(\partial \mathbf{X}_0)$ (where $\partial \mathbf{X}_0 = D(\mathbf{X}_0)_1$), satisfying the Leibniz rule with respect to the tensor product (and additional conditions). The condition that the restriction to ω is the identity corresponds to the derivation being relative to the canonical inclusion of $\omega(\mathbf{X}_0)$ in $\omega(\partial \mathbf{X}_0)$ given by the differential structure.

Similarly, a differential automorphism of ω is an automorphism t of ω as a tensor functor, with the additional condition that for any object $\mathbf{X}_0$, the diagram

$$\begin{array}{ccc} \omega(\mathbf{X}_0) & \xrightarrow{d_{\mathbf{X}_0}} & \omega(\partial \mathbf{X}_0) \\ \downarrow t_{\mathbf{X}_0} & & \downarrow t_{\partial \mathbf{X}_0} \\ \omega(\mathbf{X}_0) & \xrightarrow{d_{\mathbf{X}_0}} & \omega(\partial \mathbf{X}_0) \end{array} \quad (5)$$

commutes. Thus the condition (4) really is about preservation of the differentiation.

5.2.8. More generally, we define a *derivation* on an object $\mathbf{X}_0$ of $\mathcal{C}$ to be a morphism $d : \mathbf{X}_1 := \partial(\mathbf{X}_0) \rightarrow \mathbf{X}_0$ such that the composition $\mathbf{X}_0 \rightarrow \mathbf{X}_1 \xrightarrow{d} \mathbf{X}_0$ is the identity.

Given two derivations $d_X : \mathbf{X}_1 \rightarrow \mathbf{X}_0$ and $d_Y : \mathbf{Y}_1 \rightarrow \mathbf{Y}_0$, we define the combined derivation $d = d_X \otimes d_Y : (\mathbf{X} \otimes \mathbf{Y})_1 = (\mathbf{X}_0 \otimes \mathbf{Y}_0)_1 \rightarrow \mathbf{X}_0 \otimes \mathbf{Y}_0$ to be the restriction of $d_X \otimes Id \oplus Id \otimes d_Y$ to $(\mathbf{X} \otimes \mathbf{Y})_1$ (this makes sense, since both are the identity on $\mathbf{X}_0 \otimes \mathbf{Y}_0$). In $\mathcal{V}_{ec\mathbb{k}}$ this corresponds to a real derivation, in the sense that it gives a map $d_0 : \mathbf{X}_0 \otimes \mathbf{Y}_0 \rightarrow \mathbf{X}_0 \otimes \mathbf{Y}_0$ which is a derivation over $\mathbb{k}$ and also $d_0(x \otimes y) = d_X(x) \otimes y + x \otimes d_Y(y)$.

There is a canonical derivation d_1 on $\mathbf{1}_0$ given by the first projection $\mathbf{1}_1 = \mathbf{1}_0 \oplus \mathbf{1}_0 \rightarrow \mathbf{1}_0$. It has the property that $d_1 \otimes d = d \otimes d_1$ for any derivation d on any $\mathbf{X}_0$, under the canonical identification of $\mathbf{1} \otimes \mathbf{X}_0$ and $\mathbf{X}_0 \otimes \mathbf{1}$ with $\mathbf{X}_0$.

5.3. Differential algebraic group.

5.3.1. Let K be a differential field (i.e., a field endowed with a derivation). We recall (Kolchin [16]) that a *Kolchin closed* subset (of a affine n -space) is given by a collection of polynomial (ordinary) differential equations in variables $x_1, \dots, x_n$, i.e., polynomial equations in variables $\delta^i(x_j)$, for $i \geq 0$. Such a collection determines a set of points (solutions) in any differential field extension of K . As with algebraic varieties, it is possible to study these sets by considering points in a fixed field, provided that it is *differentially closed*. The Kolchin closed sets form a basis of closed sets for a Noetherian topology. Morphisms are also given by differential polynomials. A differential algebraic group is a group object in this category.

More generally, it is possible to consider a differential K -algebra, i.e., a K -algebra with a vector field extending the derivation on K , and develop consider these notions there.

5.3.2. By a *linear* differential algebraic group, we mean a differential algebraic group which is represented by a differential Hopf algebra. A differential algebraic group which is affine as a differential algebraic variety need not be linear in this sense, since a morphism of affine differential varieties need not correspond to a map of differential algebras. Any linear differential algebraic group has a faithful representation. All these results appear in Cassidy [2], along with an example of an affine non-linear group. In Cassidy [3] it is shown that any representation of a *linear* group (and more generally, any morphism of linear groups) does correspond to a map of differential algebras.

5.3.3. *Differential representations.* Let $\mathcal{G}$ be a linear differential algebraic group over a differential field $\mathbb{k}$. A representation of $\mathcal{G}$ is given by a finite dimensional vector space V over $\mathbb{k}$, together with a morphism $\mathcal{G} \rightarrow GL(V)$. A map of representations is a linear transformation that gives a map of group representations for each differential $\mathbb{k}$ -algebra. The category of all such representations is denoted $\mathcal{Rep}_{\mathcal{G}}$.

We endow $\mathcal{Rep}_{\mathcal{G}}$ with a differential structure in the same way as for vector spaces. If V is a representation of $\mathcal{G}$, assigning gv to (g, v) , then the action of $\mathcal{G}$ on $\mathfrak{D} \hat{\otimes} V$ is given by $(g, x \hat{\otimes} v) \mapsto x \hat{\otimes} gv$. With this differential structure, the forgetful functor ω into $\mathcal{V}_{ec\mathbb{k}}$ has an obvious structure of a differential tensor functor.

A differential automorphism t of ω is given by a collection of vector space automorphisms t_V , for any representation V of G . The commutativity condition (5) above translates to the condition that $t_{\mathfrak{D} \hat{\otimes} V} = 1 \hat{\otimes} t_V$.

In particular, given a differential $\mathbb{k}$ -algebra A , and $g \in \mathcal{G}(A)$, action by g gives an automorphism of $A \otimes \omega$ as a differential tensor functor, since the action of g on $\mathfrak{D} \hat{\otimes} V$ is deduced from its action on V . Thus we get a map $\mathcal{G} \rightarrow \mathcal{G}_\omega$. We shall prove in 5.5.5 that the map is an isomorphism.

5.3.4. Example. Let $\mathcal{G}_m$ be the (differential) multiplicative group, and let $\overline{\mathcal{G}_m}$ be the multiplicative group of the constants (thus, as differential varieties, $\mathcal{G}_m$ is given by the equation $xy = 1$, and $\overline{\mathcal{G}_m}$ is the subvariety given by $x' = 0$.) There is a differential algebraic group homomorphism dlog from $\mathcal{G}_m$ to $\mathcal{G}_a$ (the additive group), sending x to x'/x , and $x \mapsto x'$ is a differential algebraic group endomorphism of $\mathcal{G}_a$. Let V be the standard 2-dimensional algebraic representation of $\mathcal{G}_a$. Using dlog and the derivative, we get for any $i \geq 0$ a 2-dimensional irreducible representation V_i of $\mathcal{G}_m$, which are all unrelated in terms of the tensor structure (and unrelated with the non-trivial 1-dimensional algebraic representations of $\mathcal{G}_m$.)

However, if $\mathbf{X}$ is the $\mathcal{G}_m$ representation corresponding to the identity map on $\mathcal{G}_m$, an easy calculation shows that V_0 is isomorphic to $\partial \mathbf{X} \otimes \check{\mathbf{X}}$. Similarly, V_{i+1} is a quotient of ∂V_i .

The inclusion of $\overline{\mathcal{G}_m}$ in $\mathcal{G}_m$ gives a functor from $\mathcal{R}ep_{\mathcal{G}_m}$ to $\mathcal{R}ep_{\overline{\mathcal{G}_m}}$. But in $\mathcal{R}ep_{\overline{\mathcal{G}_m}}$, V_0 is isomorphic to $\mathbf{1} \oplus \mathbf{1}$ (and $\partial \mathbf{X}$ to $\mathbf{X} \oplus \mathbf{X}$.)

5.4. Differential schemes in $\mathcal{C}$.

5.4.1. We recall that the $\otimes$ operation on $\mathcal{C}$ extends canonically to $\text{Ind}(\mathcal{C})$, making it again an abelian monoidal category. The prolongation $\mathcal{P}(\text{Ind}(\mathcal{C}))$ can be identified with $\text{Ind}(\mathcal{P}(\mathcal{C}))$, and a differential structure on $\mathcal{C}$ thus extends canonically to a differential structure on $\text{Ind}(\mathcal{C})$.

Recall (Deligne [4, p. 7.5]) that if $\mathcal{C}$ is a tensor category, a *ring* in $\text{Ind}(\mathcal{C})$ is an object $\mathbf{A}$ of $\text{Ind}(\mathcal{C})$ together with maps $m : \mathbf{A} \otimes \mathbf{A} \rightarrow \mathbf{A}$ and $u : \mathbf{1} \rightarrow \mathbf{A}$ satisfying the usual axioms.

5.4.2. Definition. Let $(\mathcal{C}, D)$ be a differential tensor category, and let $\mathbf{A}$ be a commutative ring in $\text{Ind}(\mathcal{C})$. A *vector field* on $\mathbf{A}$ is a derivation on $\mathbf{A}$ in the sense of 5.2.8, which commutes with the product, and which restricts to the canonical derivation d_1 on $\mathbf{1}$. A *differential algebra* in $\text{Ind}(\mathcal{C})$ is a commutative ring in $\text{Ind}(\mathcal{C})$ together with a vector field. An (affine) *differential scheme* in $\mathcal{C}$ is a differential algebra in $\text{Ind}(\mathcal{C})$, viewed as an object in the opposite category.

5.4.3. Higher derivations. Let $\mathbf{X}_0$ be an object of a differential tensor category $(\mathcal{C}, D)$. As explained above, $D(\mathbf{X}_0)$ can be viewed as representing a universal derivation on $\mathbf{X}_0$. We now construct the analogue of higher derivatives. More precisely, we define, by induction for each $n \geq 0$, the following data:

- (1) An object $\mathbf{X}_n$ (of $\mathcal{C}$)
- (2) A map $q_n : \mathbf{X}_{n-1} \rightarrow \mathbf{X}_n$
- (3) A map $t_n : \partial \mathbf{X}_{n-1} \rightarrow \mathbf{X}_n$

such that $q_{n+1} \circ t_n = t_{n+1} \circ \partial(q_n)$. In the context of $\mathcal{V}ec_{\mathbb{k}}$, the data can be thought of as follows: $\mathbf{X}_n$ is the space of expressions $v_0 + \partial v_1 + \cdots + \partial^n v_n$ with $v_j \in \mathbf{X}_0$; q_n the inclusion of elements as above; the map t_n linear map corresponding to the derivation.

For the base, setting $\mathbf{X}_{-1} = 0$ determines all the data in an obvious way. Given $\mathbf{X}_n$, $t_{n+1} : \partial(\mathbf{X}_n) \rightarrow \mathbf{X}_{n+1}$ is defined to be the co-equaliser of the following two maps:

$$\begin{array}{ccc} & \mathbf{X}_n & \\ t_n \nearrow & & \searrow i \\ \partial(\mathbf{X}_{n-1}) & \xrightarrow{\hat{\partial}(q_n)} & \partial(\mathbf{X}_n) \end{array} \quad (6)$$

Where i is part of the structure of $D(\mathbf{X}_n)$. The map q_{n+1} is the composition $t_{n+1} \circ i$. Clearly the commutativity condition is satisfied. We note that the two objects we denote by $\mathbf{X}_1$ coincide, and the map q_1 coincides with the map i for $D(\mathbf{X}_0)$. The map t_1 is the identity.

5.4.4. Definition. Let $\mathbf{X}_0$ be an object of $\mathcal{C}$. The *differential scheme associated with $\mathbf{X}_0$* , denoted $\mathbb{A}(\mathbf{X}_0)$, is a differential scheme in $\mathcal{C}$ defined as follows: Let $\mathbf{D}$ be the ind-object defined by the system $\tilde{\mathbf{X}}_i$, with maps q_i (as in 5.4.3). The maps t_i there define a derivation t on $\mathbf{D}$. This derivation induces a derivation on tensor powers of $\mathbf{D}$ (as in 5.2.8), which descends to the symmetric powers. It is easy to see that this determines a differential algebra structure on the symmetric algebra on $\mathbf{D}$. $\mathbb{A}(\mathbf{X}_0)$ is the associated scheme.

A morphism in $\mathcal{C}$ clearly determines a morphism of schemes on the associated schemes, making $\mathbb{A}(-)$ into a functor.

5.5. Model theory of differential fibre functors. We now wish to prove statements analogous to the ones for algebraic Tannakian categories, using the same approach as in section 4. We work with a fixed differential tensor category $(\mathcal{C}, D)$, with $\mathbb{k} = \text{End}(\mathbf{1})$ a field. We view $\mathbb{k}$ as a differential field, with the differential structure induced from D , as in 5.2.2.

We will be using the theory DCF of differentially closed fields. We refer the reader to Marker [20] or Marker et al. [21] for more information.

5.5.1. The theory associated with a fibre functor. The theory $T_{\mathcal{C}}$ associated with the data above, as well as a differential field extension K of $\mathbb{k}$ is an expansion of the theory $T_{\mathcal{C}}$ as defined in 4.1 by the following structure:

- (1) $\mathbf{L}$ has an additional function symbol $'$, and the theory says that $'$ is a derivation, and that $\mathbf{L}$ is a differentially closed field (and with the restriction of $'$ to K as given).
- (2) For every object $\mathbf{X}$, there is a function symbol $d_{\mathbf{X}} : V_{\mathbf{X}} \rightarrow V_{\partial(\mathbf{X})}$. This function is a derivation, in the sense that for any $a \in \mathbf{L}$ and $v \in V_{\mathbf{X}}$,

$$d_{\mathbf{X}}(av) = a'V_{i_{\mathbf{X}}}(v) + ad_{\mathbf{X}}(v)$$

The theory furthermore says that $d_{\mathbf{X}}$ identifies $V_{\partial(\mathbf{X})}$ with $\mathfrak{D} \hat{\otimes} V_{\mathbf{X}}$ (in any model), in the sense of 5.2.7 (explicitly, it says that $V_{p_{\mathbf{X}}} \circ d_{\mathbf{X}}$ is the identity map.)

- (3) The maps d and b (from the tensor structure) are compatible with the structure of tensor functor of D : given objects $\mathbf{X}$ and $\mathbf{Y}$ of $\mathcal{C}$, let $c_{\mathbf{X}, \mathbf{Y}} : \partial(\mathbf{X} \otimes \mathbf{Y}) \rightarrow (\partial(\mathbf{X}) \otimes \partial(\mathbf{Y}))_1$ be the isomorphism supplied with D . Then we require that $V_{c_{\mathbf{X}, \mathbf{Y}}} \circ d_{\mathbf{X} \otimes \mathbf{Y}} \circ b_{\mathbf{X}, \mathbf{Y}}$ coincides with $b_{\partial(\mathbf{X}), \mathbf{Y}} \circ d_{\mathbf{X}} \times 1 + b_{\mathbf{X}, \partial(\mathbf{Y})} \circ 1 \times d_{\mathbf{Y}}$.

5.5.2. Let ω be a differential fibre functor on $\mathcal{C}$, and let $L = M_{\mathbf{1}}$ be a differentially closed field containing K . As in section 4, we expand $M_{\mathbf{1}}$ to a model M of $T_{\mathcal{C}}$ by tensoring with L . The differential structure of ω gives (as in 5.2.7) a universal derivation $\omega(\mathbf{X}) \rightarrow \omega(\partial(\mathbf{X}))$, which extends uniquely to a (universal) derivation $(d_{\mathbf{X}})_M$ on $M_{\mathbf{X}}$.

As in the algebraic case, the model M_{ω} just constructed contains, in a natural way, the subset of elements coming from the fibre functor ω . This set, which will also be denoted by ω is definably closed, as before. In particular, we get, as before, the following corollary (with $K = \mathbb{k}$):

Corollary. *Assume that $\mathcal{C}$ has a differential fibre functor over K . Then $T_{\mathcal{C}}$ is consistent, and (in a model) $\text{dcl}(0) \cap \mathbf{L} = \mathbb{k}$.*

5.5.3. *Internality.* Since the differential $T_{\mathcal{C}}$ is an expansion of the algebraic one with no new sorts, it is again an internal cover of $\mathbf{L}$. Furthermore, if B is a basis for some $V_{\mathbf{X}}$, then $B \cup d_{\mathbf{X}}(B)$ is a basis for $V_{\partial(\mathbf{X})}$. Therefore, if $\mathcal{C}$ is generated as a differential tensor category by one object (in the sense that the objects $\partial^i \mathbf{X}$ generate $\mathcal{C}$ as a tensor category), then all of the sorts are internal using the same finite parameter. As usual, the general case is obtained by taking a limit of such.

5.5.4. Let ω be a fibre functor, and recall (5.2.6) that G_{ω} denotes the group functor $A \mapsto \underline{\text{Aut}}^{\circ}(A \otimes \omega)$ on the category of differential $\mathbb{k}$ -algebras. As in the algebraic case, G_{ω} is isomorphic to $A \mapsto G(A \otimes_{\mathbb{k}} \omega)$, where G is the (pro-) definable group associated with the internal cover $T_{\mathcal{C}}$ (and A is any differential field extension of $\mathbb{k}$).

5.5.5. Let H be a differential algebraic group over $\mathbb{k}$, let $\mathcal{C} = \mathcal{R}ep_H$ (example 5.3.3), and let ω be the forgetful functor. We consider H as a functor on differential field extensions of $\mathbb{k}$. As before, the action of H on its representations gives rise to a definable, and therefore differential algebraic map $H \rightarrow G_{\omega} = G(- \otimes \omega)$. As in the algebraic case, we would like to conclude that this is a bijection. For this purpose, we need to determine the imaginaries in $T_{\mathcal{C}}$. The following is an analogue of Hrushovski [12, Proposition 4.2]. The rest of the proof is precisely as before.

5.5.6. *Proposition.* $T_{\mathcal{C}}$ eliminates imaginaries to the level of projective spaces.

Proof. We need to show that any definable set S over parameters can be defined with a canonical parameter. Since, by assumption, no new structure is induced on $\mathbf{L}$, and any set is internal to $\mathbf{L}$, every such set is Kolchin constructible. By Noetherian induction, it is enough to consider S Kolchin closed.

We claim that the algebra of differential polynomials on a sort U is ind-definable in $T_{\mathcal{C}}$. Indeed, it is precisely given by the interpretation of scheme structure associated with U (5.4.4). We only mention the definition of the evaluation map (using the notation there): it is enough to the evaluation on (the interpretation of) $\mathbf{D}$, since the rest is as in the algebraic case. We define the evaluation $e_n : \check{U}_n \times U \rightarrow \mathbf{L}$ by induction on n : e_0 is the usual evaluation. If $u \in U$, the map $d \mapsto e_n(d, u)'$ is a derivation on $\check{U}_n$, and so defines a linear map from $\partial(\check{U}_n)$ to $\mathbf{L}$. Inspection of the definition of $\check{U}_{n+1}$ (for vector spaces) reveals that this map descends to a linear map $e_{n+1}(-, u)$ on $\check{U}_{n+1}$.

The rest of the proof is the same as in Hrushovski [12], namely, the Kolchin closed set S is determined by the finite dimensional linear space spanned by the

defining equations, and this space is an elements of some Grassmanian, which is, in turn, a closed subset of some projective space. $\square$

5.5.7. We note that conversely, given $\mathcal{C}$ and ω , the automorphism group G is a Kolchin closed subgroup of $GL(V_{\mathbf{X}})$, where $\mathbf{X}$ generates $\mathcal{C}$ as a differential tensor category, and the category of definable $\mathbf{L}$ vector spaces in $T_{\mathcal{C}}$ and definable linear maps between them is equivalent to $\mathcal{R}ep_G$.

REFERENCES

- [1] Michael Artin and Barry Mazur. *Etale homotopy*. Lecture Notes in Mathematics 100. Reprint of the 1969 original. Berlin: Springer-Verlag, 1986. ISBN: 3-540-04619-4. See p. 5.
- [2] Phyllis Joan Cassidy. “Differential algebraic groups”. In: *Amer. J. Math.* 94 (1972), pp. 891–954. ISSN: 0002-9327. See p. 25.
- [3] Phyllis Joan Cassidy. “The differential rational representation algebra on a linear differential algebraic group”. In: *J. Algebra* 37.2 (1975), pp. 223–238. ISSN: 0021-8693. See p. 25.
- [4] Pierre Deligne. “Catégories tannakiennes”. In: *The Grothendieck Festschrift*. Vol. II. Progress in Mathematics 87. Boston, MA: Birkhäuser Boston Inc., 1990, pp. 111–195. ISBN: 0-8176-3428-2. See pp. 1, 2, 19, 26.
- [5] Pierre Deligne and James S. Milne. “Tannakian categories”. In: *Hodge cycles, motives, and Shimura varieties*. Lecture Notes in Mathematics 900. Berlin: Springer-Verlag, 1982. Chap. II, pp. 101–228. ISBN: 3-540-11174-3. URL: <http://jmilne.org/math>. See pp. 1–3, 5, 14, 19–21.
- [6] Peter Freyd. *Abelian categories. An introduction to the theory of functors*. Harper’s Series in Modern Mathematics. New York: Harper & Row Publishers, 1964. See p. 21.
- [7] Dennis Gaitsgory and David Kazhdan. “Representations of algebraic groups over a 2-dimensional local field”. In: *Geom. Funct. Anal.* 14.3 (2004), pp. 535–574. ISSN: 1016-443X. arXiv: [math/0302174](https://arxiv.org/abs/math/0302174). See p. 5.
- [8] Alexander Grothendieck et al. *Revêtements étales et groupe fondamental (SGA 1)*. Documents Mathématiques (Paris) 3. Séminaire de géométrie algébrique du Bois Marie 1960–61., Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)]. Paris: Société Mathématique de France, 2003. ISBN: 2-85629-141-4. arXiv: [math/0206203](https://arxiv.org/abs/math/0206203). See p. 20.
- [9] Alexander Grothendieck et al. *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 269. Berlin: Springer-Verlag, 1972. See p. 5.
- [10] Bradd Hart and Ziv Shami. “On the type-definability of the binding group in simple theories”. In: *J. Symbolic Logic* 70.2 (2005), pp. 379–388. ISSN: 0022-4812. See p. 2.

- [11] Ehud Hrushovski. “Computing the Galois group of a linear differential equation”. In: *Differential Galois theory*. Ed. by Teresa Crespo and Zbigniew Hajto. Banach Center Publications 58. May 28–June 1, 2001. Warsaw: Polish Academy of Sciences Institute of Mathematics, 2002, pp. 97–138. See pp. 2, 10.
- [12] Ehud Hrushovski. *Groupoids, imaginaries and internal covers*. Mar. 2006. arXiv: [math/0603413](https://arxiv.org/abs/math/0603413). See pp. 2, 8, 10, 16, 19, 28, 29.
- [13] Ehud Hrushovski. “Unidimensional theories are superstable”. In: *Ann. Pure Appl. Logic* 50.2 (1990), pp. 117–138. ISSN: 0168-0072. See p. 2.
- [14] Moshe Kamensky. “Definable groups of partial automorphisms”. Accepted to *Selecta Mathematica*. July 2006. See p. 14.
- [15] Moshe Kamensky. “Ind- and Pro- definable sets”. In: *Ann. Pure Appl. Logic* 147.3 (2007), pp. 180–186. ISSN: 0168-0072. DOI: [10.1016/j.apal.2007.04.003](https://doi.org/10.1016/j.apal.2007.04.003). arXiv: [math/0608163](https://arxiv.org/abs/math/0608163). See p. 5.
- [16] Ellis R. Kolchin. *Differential algebra and algebraic groups*. Pure and Applied Mathematics 54. New York: Academic Press, 1973. See p. 25.
- [17] Saunders Mac Lane. *Categories for the working mathematician*. 2nd ed. Graduate Texts in Mathematics 5. New York: Springer-Verlag, 1998. ISBN: 0-387-98403-8. See pp. 3, 4.
- [18] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in geometry and logic*. Universitext. New York: Springer-Verlag, 1994. ISBN: 0-387-97710-4. See pp. 5, 6.
- [19] Michael Makkai and Gonzalo E. Reyes. *First order categorical logic*. Lecture Notes in Mathematics, Vol. 611. Model-theoretical methods in the theory of topoi and related categories. Berlin: Springer-Verlag, 1977. ISBN: 3-540-08439-8. See pp. 11, 12.
- [20] David Marker. “Model theory of differential fields”. In: *Model theory, algebra, and geometry*. Ed. by Deirdre Haskell, Anand Pillay, and Charles Steinhorn. Mathematical Sciences Research Institute Publications 39. Cambridge: Cambridge University Press, 2000, pp. 53–63. ISBN: 0-521-78068-3. URL: <http://www.msri.org/communications/books/Book39/contents.html>. See p. 27.
- [21] David Marker, Margit Messmer, and Anand Pillay. *Model theory of fields*. Second. Lecture Notes in Logic 5. La Jolla, CA: Association for Symbolic Logic, 2006. ISBN: 978-1-56881-282-3; 1-56881-282-5. See p. 27.
- [22] James S. Milne. *Étale cohomology*. Princeton Mathematical Series 33. Princeton, N.J.: Princeton University Press, 1980. ISBN: 0-691-08238-3. See p. 17.
- [23] Alexey Ovchinnikov. “Tannakian approach to linear differential algebraic groups”. In: *Transform. Groups* 13.2 (2008), pp. 413–446. ISSN: 1083-4362. See p. 2.
- [24] Alexey Ovchinnikov. “Tannakian categories, linear differential algebraic groups, and parametrized linear differential equations”. In: *Transform. Groups* 14.1 (2009), pp. 195–223. ISSN: 1083-4362. See p. 2.
- [25] Anand Pillay. “Algebraic D -groups and differential Galois theory”. In: *Pacific J. Math.* 216.2 (2004), pp. 343–360. ISSN: 0030-8730. See p. 2.
- [26] Bruno Poizat. *Stable groups*. Mathematical Surveys and Monographs 87. Providence, RI: American Mathematical Society, 2001. ISBN: 0-8218-2685-9. See p. 18.

- [27] Bruno Poizat. “Une théorie de Galois imaginaire”. In: *J. Symbolic Logic* 48.4 (1983), 1151–1170 (1984). ISSN: 0022-4812. See p. 1.
- [28] Neantro Saavedra-Rivano. *Catégories Tannakiennes*. Lecture Notes in Mathematics, Vol. 265. Berlin: Springer-Verlag, 1972. See pp. 1, 14.
- [29] Gerald E. Sacks. *Saturated model theory*. Reading, Mass.: W. A. Benjamin, Inc., 1972. See p. 9.
- [30] William C. Waterhouse. *Introduction to affine group schemes*. Graduate Texts in Mathematics 66. New York: Springer-Verlag, 1979. ISBN: 0-387-90421-2. See p. 18.
- [31] Boris I. Zil’ber. “Totally categorical theories: structural properties and the nonfinite axiomatizability”. In: *Model theory of algebra and arithmetic*. Karpacz, September 1–7, 1979. Ed. by L. Pacholski and J. Wierzejewski. Lecture Notes in Mathematics 834. Berlin: Springer, 1980, pp. 381–410. ISBN: 3-540-10269-8. See p. 1.

DEPARTMENT OF PURE MATH, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA

E-mail address: <mailto:mkamensky@math.uwaterloo.ca>

URL: <http://mkamensky.notlong.com>